

Tweed River Entrance Sand Bypassing Project

Kirra Reef Marine Biota Monitoring 2014

Prepared for:

NSW Department of Trade and Investment, Crown Lands

frc environmental

PO Box 2363, Wellington Point QLD 4160

Telephone: + 61 3286 3850 Facsimile: + 61 3821 7936

frc reference: 131218





Document Control Summary

Project No.: 131218
Status: Draft Report
Project Director: John Thorogood
Project Manager: Craig Chargulaf

Title: TRESBP Kirra Reef Marine Biota Monitoring 2014

Project Team: C. Chargulaf, M. Langridge, A. Lea, L. Thorburn, S. Walker.

Client: Department of Trade and Investment, Crown Lands

Client Contact: Catherine Kerr

Date: May 2014

Edition: 131218Ri

Checked by: Lauren Thorburn _____

Issued by: Craig Chargulaf _____

Distribution Record

Trade and Investment, Crown Lands: 2 copies (1 Word, 1 PDF)

This work is copyright.

A person using frc environmental documents or data accepts the risk of:

- 1 Using the documents or data in electronic form without requesting and checking them for accuracy against the original signed hard copy version; and
- 2 Using the documents or data for any purpose not agreed to in writing by frc environmental.

Contents

Sum	mary		i					
1	Intro	duction	1					
	1.1	Historical Context of Monitoring	1					
	1.2	Sand Nourishment History	2					
	1.3	Faunal and Floral Characteristics of the Survey Region	7					
2	Meth	nods	9					
	2.1	Benthic Flora and Macroinvertebrates	9					
	2.2	Fish	14					
	2.3	B Data Analysis						
3	Resu	ults	16					
	3.1	Cover of Benthic Assemblages	16					
	3.2	Benthic Algae	22					
	3.3	Benthic Macro-Invertebrates	26					
	3.4	Fish	36					
4	Disc	ussion	39					
	4.1	Changes to the Ecological Condition of Kirra Reef	39					
	4.2	Impacts of the Sand Bypassing System on Kirra Reef	43					
	4.3	Impacts of Storms & Seasonality on Kirra Reef	44					
	4.4	Long-term Impacts of the Sand Bypassing System on Kirra Reef	45					
5	Cond	clusions	46					
6	Refe	rences	48					
Appe	endix A	A Introduction to Data Analysis Used.						
Арре	endix E	Cover and Abundance of Benthic Fauna and Flora on Kirra Beach Reefs in April 2014.	and Palm					

Appendix C Relative Abundance of Fish Found at Kirra and Palm Beach Reefs in Each Survey.

Tables

Table 3.1	PERMANOVA results for multivariate differences in the composition of benthic assemblages between surveys and	
	locations.	16
Table 3.2	Pairwise ANOSIM results for differences in assemblage composition between reefs for each survey.	16
Table 3.3	PERMANOVA post hoc pairwise results for the differences between locations in each survey events, for the assemblage composition and cover of selected benthic groups.	17
Table 3.4	Univariate PERMANOVA results for differences in the cover of macroalgae between surveys and location.	24
Table 3.5	Univariate PERMANOVA results for differences in the cover of turf algae between surveys and locations.	26
Table 3.6	Univariate PERMANOVA results for differences in the cover of sponges between surveys and locations.	27
Table 3.7	Univariate PERMANOVA results for the differences in cover of all ascidians other than <i>Pyura stolonifera</i> between surveys and locations.	30
Table 3.8	Univariate PERMANOVA results for the differences in the cover of hard coral between surveys and locations.	32
Table 3.9	Univariate PERMANOVA results for the differences in the cover of soft coral between surveys and locations.	33
Table 3.10	Univariate PERMANOVA results for the differences in crinoid density between surveys and locations.	35

Figures

Figure 1.1	Extent of Kirra Reef in 2001, 2003 and 2004.	4
Figure 1.2	Extent of Kirra Reef in 1995, 2001, 2011 and 2012.	5
Figure 1.3	Extent of Kirra Reef in 2014.	6
Figure 2.1	Location of (a) Palm Beach Reef and (b) Kirra Reef	9
Figure 2.2	Small exposed outcrop in the eastern section of Kirra Reef.	10
Figure 2.3	Diver surveying benthic assemblages at Kirra Reef in April 2014.	10
Figure 3.1	Multi-dimensional scaling plot of benthic cover in the 1995, 1996, 2001 and 2003 surveys.	18
Figure 3.2	Multi-dimensional scale plot of benthic cover in the 2004, 2005, 2010 and 2012 surveys.	19
Figure 3.3	Multi-dimensional scale plot of benthic cover in the 2014 survey.	20
Figure 3.4	Multi-dimensional scale plot of benthic cover at Kirra Reef in April 1995 and April 2014.	21
Figure 3.5	Sargassum sp. dominated the macroalgal communities of Kirra Reef at site KN3 in April 2014.	23
Figure 3.6	Mean cover of macroalgae (\pm SE) at Kirra Reef and Palm Beach Reef in all surveys.	23
Figure 3.7	Areas of Palm Beach Reef support little macroalgae but have a high cover of turf algae (April 2014).	25
Figure 3.8	Mean cover of turf algae (\pm SE) at Kirra Reef and Palm Beach Reef in all surveys.	25
Figure 3.9	Sponges were much more abundant at Palm Beach Reef than at Kirra Reef since 2005.	26
Figure 3.10	Mean cover of sponges (\pm SE) at Kirra Reef and Palm Beach Reef on all surveys.	27
Figure 3.11	Ascidians (<i>Pyura stolonifera</i> and <i>Cnemidocarpa stolonifera</i>) were present at Kirra Reef in April 2014.	28

Figure 3.12	Mean abundance of ascidians (<i>Pyura stolonifera</i>) (individuals / $0.25~\text{m}^2$) (\pm SE) at Kirra Reef and Palm Beach Reef in all surveys.	29
Figure 3.13	Mean cover of ascidians (\pm SE), other than <i>Pyura stolonifera</i> , at Kirra Reef and Palm Beach Reef in all surveys.	30
Figure 3.14	Mean cover of hard corals (\pm SE) at Kirra Reef and Palm Beach Reef in all surveys.	31
Figure 3.15	Mean cover of soft corals (\pm SE) at Kirra Reef and Palm Beach Reef in all surveys.	33
Figure 3.16	Crinoids (feather stars) were patchily distributed found at Kirra and Palm Beach reefs in April 2014.	34
Figure 3.17	Mean abundance of crinoids (individuals / 0.25 m 2) (\pm SE) at Kirra Reef and Palm Beach Reef in all surveys.	35
Figure 3.18	Number of fish species recorded at Kirra Reef and Palm Beach Reef on each survey.	37
Figure 3.19	Yellowtail were abundant at Kirra Reef in April 2014.	37
Figure 3.20	Barred soapfish were recorded at Palm Beach Reef for the first time in April 2014.	38
Figure 3.21	Cowtail stingray was recorded at Kirra Reef for the first time in April 2014.	38
Figure 3.22	Large scale parma continued to be abundant at Palm Beach Reef in April 2014.	38
Maps		
Мар 1	Sites surveyed at Kirra Reef in 2014.	12
Мар 2	Sites surveyed at Palm Beach Reef in 2014.	13

Summary

The NSW Department of Trade and Investment, Crown Lands commissioned frc environmental on behalf of the Tweed River Entrance Sand Bypassing Project (TRESBP) to monitor the condition and biodiversity of benthic assemblages at Kirra Reef, and to assess the impacts of the project on those assemblages. The purpose of the TRESBP is to maintain a navigable entrance to the Tweed River, and to provide a continuing supply of sand to the southern Gold Coast beaches that is consistent with the natural rate of longshore drift. This report discusses the results of ecological monitoring of the benthic fauna, flora, and fish of Kirra Reef, completed in April 2014.

Ongoing monitoring of Kirra Reef is required under the Environmental Management System (EMS) Sub-Plan B14 Kirra Reef Management Plan, prepared by the TRESBP in February 2001. The methods used in the April 2014 survey (i.e. surveys of benthic cover and fish abundance), were those developed for the Stage I survey completed in 1995, and have been used in the subsequent surveys in 1996, 2001, 2003, 2004, 2005, 2010 and 2012.

Impacts of the Sand Bypassing System on Kirra Reef

During the early years of the TRESBP operation, large amounts of sand were deposited on the southern Gold Coast beaches. This was done to provide a 'catch up' quantity of sand to the badly eroded beaches, reduce the Tweed River entrance bar and clear a sand trap in the vicinity of the sand collection jetty to increase the efficiency of the bypassing system. During the initial period of increased deposition, the volume of sand delivered exceeded the amount that was transported north through natural sand transport mechanisms. This resulted in a large volume of sand being deposited on the southern beaches, and wave action and tidal currents redistributed some of this sand over Kirra Reef. This resulted in smothering and a decline in the areal extent of Kirra Reef. The project EIS predicted that the impact associated with the gradual accumulation of sand around the base of Kirra Reef was unavoidable.

Since the delivery of large quantities of sand was completed in 2005, the volume of sand delivered by the project has declined, and now matches the natural rate of northward sand transport more closely. However, there was a substantial lag between the reduction in sand delivery and transport of the sand further north, due to a period of calmer than usual conditions with reduced storm activity from the north-east. As such, dispersion of sand from Kirra Beach and reduction in the sand levels around the reef was slower than predicted between 2005 and 2010.

Between February 2010 and July 2012, there was a large (50%) increase in the area of exposed rock in the northern section of Kirra Reef, which is directly correlated with the migration of sand further to the north. However, between July 2012 and April 2014 there was a minor reduction in the area of exposed rock. Nonetheless, as predicted in the project EIS, the extent of the reef uncovered in the northern section now closely approximates that recorded before the Tweed River training walls were extended. Similar to previous surveys, in 2014 the eastern and inner western parts of the reef remained buried due to their close proximity to the shoreline.

Changes to the Ecological Condition of Kirra Reef

The Project's EIS did not consider the ecological consequences of both the reduced areal extent and increased wave energy (a consequence of decreased depth of the reef habitat available) that would occur as a result of the accretion of sand around the base of the rock outcrops at Kirra Reef. However, once sand transport became more consistent with the natural supply, it was expected that the benthic flora and fauna assemblages of the reef would return to a condition consistent with the historical reef extent and natural sand transport patterns (and associated coastal fluctuations, wave action, sedimentation and water quality) that were observed in the vicinity of the reef prior to the development of the Tweed River training walls.

Since monitoring began, the greatest change to the ecological assemblages on Kirra Reef has been due to the loss of large areas of rocky reef, caused by burial of the rock by sand. The loss of reef habitat has reduced the area of hard substratum available for colonisation and consequently the overall abundance (cover) of benthic sessile assemblages. The redistribution of sand also resulted in small-scale changes to the distribution and cover of the benthic assemblages on the remaining reef over time, through complex interactions between physical disturbance (i.e. increased burial, sedimentation, wave action, and abrasion), food availability, competition, and local weather and sea conditions.

In April 2014, the diversity of assemblages had increased relative to July 2012, as did the cover of macroalgae (though it remained well below the peak of 60% cover recorded in January 2001), which is most likely due to natural succession with more mature communities being established between 2012 and 2014 when the extent of exposed rock has been relatively stable. Nevertheless, the benthic assemblage on Kirra Reef exhibited signs of ongoing stress from physical disturbance such as storm and wave disturbance, physical abrasion and burial by sand; including reduced cover of hard coral and soft coral relative to times when the level of sand was much lower. Physical disturbance from sand burial, sand abrasion and the action of storm waves appear to have historically kept the benthic assemblages on Kirra Reef in a state of early succession; however, the diversity of benthic assemblages is predicted to increase if the extent of reef remains the same or

increases, and physical disturbance remains relatively benign. This was evident in 2014 with an increased cover of benthic invertebrates (sponges and ascidians) and a reduction of the cover of turf algae.

In April 2014, a more diverse assemblage of fish was found on Kirra Reef than on Palm Beach Reef. As fish are mobile, they can move more easily to areas that are less disturbed or that exhibit more suitable conditions. While the composition of the assemblage differed from previous survey events, the differences were more likely due to the effects of seasonal changes in water temperature and the effects of prevailing conditions at the time of the survey, rather than any substantial effect of the bypassing project.

Despite some impacts from the TRESBP, overall the composition of the flora and fauna assemblages on Kirra Reef was more similar to that found at nearby Palm Beach Reef than in previous years. Kirra Reef therefore continues to provide habitat to a range of flora and fauna, and provides important marine ecological functions and services in the region. It is possible that as sand levels have stabilised over the past two years, assemblages are slowly beginning to become more similar to those recorded prior to implementation of the TRESBP, and to those at nearby unaffected reefs such as Palm Beach Reef.

Impacts of Storms & Seasonality on Kirra Reef

The large quantities of sand that were initially delivered by the project caused a substantial shallowing of the nearshore area around the reef. This increase in bed levels was responsible for increasing the incidence of wave disturbance and sand scouring around the reef, which negatively impacted on the benthic fauna and flora found growing there.

As the delivery of sand through the bypassing project now matches the natural rate of northern longshore sand transport, short-term and seasonal changes in the areal extent of the reef are more likely the result of the action of storm waves and currents shifting sand offshore, than a discrete impact of the sand bypassing activity. It was noted that short-term fluctuations that result from storms or changes in the coastal sand supply, would have been a component of the natural range of ecological conditions observed prior to the development of the training walls.

The proximity of the reef to the coast means that the benthic assemblages will continue to be affected by sand abrasion, wave disturbance and sand smothering; however, having greater balance between the delivery of sand through the project and the natural movement of sand on and offshore, is likely to result in better ecological outcomes for the benthic assemblages found on Kirra Reef and greater consistency in the extent of reef habitat that is uncovered.

Long-term Impact of the Sand Bypassing System on Kirra Reef

In April 2014, Kirra Reef covered approximately 40 to 45% (2920 m²) of the area present in 1995 (i.e. prior to the operation of the TRESBP). This is largely due to the reduction in the inner western and eastern sections of the reef, which was predicted to occur in the EIS.

We expect that the area of reef uncovered will continue to change due to seasonal shifts in sand delivery and storms; however, the diversity of flora and fauna assemblages on Kirra Reef should increase gradually over time, especially if the extent of the rocky reef that remains uncovered is consistent or increases over time to become more similar with that found prior to the development of the training walls. In this scenario, it was expected that that newly exposed areas of Kirra Reef in 2012 that were dominated by turf algae, would be colonised by other organisms including macroalgae, sponges, ascidians and potentially hard and soft coral over time. The results from 2014 indicate that this process is slowly occurring on Kirra Reef.

Ongoing monitoring will provide insight into the rate of 'recovery' of communities. It is recommended that monitoring be repeated in early 2015 (between February and April, when previous surveys have been undertaken) to confirm the results of the April 2014 survey and account for seasonal / interannual differences. It is also recommended that surveys be undertaken in 2015 in order to monitor the rate of recovery of Kirra Reef. If communities surveyed in 2015 demonstrate consistent results, monitoring could be completed bi-annually to monitor community succession, unless triggered by a 15% change in the extent of the exposed reef.

1 Introduction

frc environmental was commissioned by the NSW Government Department of Trade and Investment, Crown Lands on behalf of the Tweed River Entrance Sand Bypassing Project (TRESBP) to monitor the condition, abundance and biodiversity of floral and faunal communities at Kirra Reef. This report presents results of the survey of benthic flora and macro-invertebrate fauna, and fish at sites on Kirra Reef and at comparative sites on Palm Beach Reef, in April 2014.

The current condition of Kirra Reef was compared with the condition of nearby Palm Beach Reef in April 2014, and with changes to the reef community over time with previous assessments of Kirra Reef in 1995, 1996, 2001, 2003, 2004, 2005, 2010 and 2012 (frc environmental 2012).

1.1 Historical Context of Monitoring

Kirra Beach receives indirect sand nourishment as part of the Tweed River Entrance Sand Bypassing Project (TRESBP). The purpose of the TRESBP is to maintain a navigable entrance to the Tweed River, and to provide a continuing supply of sand to the southern Gold Coast beaches that is consistent with the natural rate of longshore drift.

frc environmental completed a baseline assessment of Kirra Reef in 1995 (Fisheries Research Consultants 1995a; b), and has undertaken seven subsequent ecological monitoring surveys of the reef on behalf of TRESBP since 1996 (Fisheries Research Consultants 1996; frc environmental 2001; 2003; 2004; 2005; 2010; 2012). Kirra Reef is the collective name given to the complex of rocky outcrops located offshore of Kirra Beach, in water depths between 3 and 7 m. The ongoing monitoring of Kirra Reef meets the requirements of the Environmental Management System (EMS) Sub-Plan B14 Kirra Reef Management Plan, prepared by the TRESBP in February 2001. It also incorporates additional monitoring activities implemented by TRESBP in August 2004.

Under Sub-Plan B14, if the area of exposed reef on aerial photographs is smaller than the range of areas shown on aerial photographs from 1962 to 1965, then monitoring of the marine biota of Kirra Reef is required. Interpretation of aerial photographs taken in August 2002 indicated there had been substantial changes to the reef since 1965. Loss of reef area continued for some years, with aerial photographs from November 2003 showing the inner northern reef and entire eastern reef covered by sand (Figure 1.1 and Figure 1.2). The area of the outer northern reef was also greatly reduced (Figure 1.2). In April 2004 the extent of the outer northern reef was further reduced, although a small outcrop of eastern reef had been uncovered (Figure 1.2). By early 2006, the area of exposed reef

had been reduced to 100 m². As a consequence of the extensive burial of the reef, ecological surveys were postponed between 2006 and 2010. Between 2006 and 2010, visual inspections of the reef were completed in place of full ecological surveys, by dive teams from Gilbert Diving and Gold Coast City Council.

By February 2010, the area of Kirra Reef had increased again and ecological surveys recommenced. In July 2012, analysis of aerial photographs indicated that the exposed section of Kirra Reef has increased in extent since the previous survey in 2010; although, the reef was still 50% smaller than recorded in 1962 and 1995 before the sand bypassing project began (Figure 1.2) (Boswood & Murray 1997). In April 2014, analysis of aerial photographs indicated a small decline in reef area since 2012 and a survey was commissioned to assess changes in the ecological communities (Figure 1.3).

1.2 Sand Nourishment History

Initial sand nourishment works (Stage 1 of the TRESBP) involved two sub-stages: Stage 1A from April 1995 to August 1996, and Stage 1B from September 1997 to May 1998. These sub-stages delivered about three million cubic metres (m³) of clean marine sand (with less than 3% fines) out to 10 metres mean water depth, including approximately 600,000 m³ of clean marine sand placed on the upper beaches during Stage 1A activities.

Prior to the establishment of the permanent sand bypassing system, additional dredging activities were undertaken to maintain a clear navigation channel at the Tweed River entrance, resulting in approximately 480,000 m³ of clean marine sand being placed in nearshore areas from Point Danger to Coolangatta Beach (from April 2000 to February 2001).

Operation of the TRESBP commenced in May 2001. Between 2001 and 2012, approximately 5.9 million m³ of pumped sand and 1.4 million m³ of dredged sand (derived from dredging of the Tweed River mouth) was deposited along the southern Gold Coast beaches. Most of the sand delivered through pumping and dredging was placed in the primary placement area, south east of Snapper Rocks. In 2013, a total of 319,883 m³ was pumped and in the current contract year, which ended in April 2014, an additional 236,278 m³ of sand was pumped to this area. Wave and current action has transported much of the sand around Snapper Rocks, nourishing beaches further to the north. Sand is also discharged from outlets at Duranbah Beach and occasionally at Snapper Rocks West. There is an outlet at Kirra; however, this has not been used since December 2003. Accumulation of sand on Kirra Reef has occurred through sand transport by waves and currents along the seabed, rather than by direct depositional smothering (Hyder Consulting Pty Ltd et al. 1997).

Relatively high quantities of sand were delivered to the southern Gold Coast beaches during the early operational years of the TRESBP to:

- provide much needed sand nourishment to the severely eroded southern Gold Coast beaches
- · reduce the Tweed Entrance Bar, and
- clear a sand trap in the vicinity of the jetty to improve the efficiently of the sand bypass system.

These project objectives were achieved, and the quantity of sand delivered since 2005 has been more consistent with the natural quantity of sand movement along the coast.

hern Reef



Figure 1.1 Extent of Kirra Reef in 2001, 2003 and 2004.




Figure 1.2 Extent of Kirra Reef in 1995, 2001, 2011 and 2012.



Figure 1.3 Extent of Kirra Reef in 2014.

1.3 Faunal and Floral Characteristics of the Survey Region

The subtidal rocky reefs of the Gold Coast region comprise remnants of highly eroded volcanic substratum that are isolated from each other by wide, variable expanses of soft sediment (Edwards & Smith 2005). They support assemblages of benthic fauna and flora, and fish that are indicative of a transition between the tropical waters of the Great Barrier Reef and the temperate waters characteristic of the mid-New South Wales coast (Done 1982; Cannon et al. 1987). These communities are dominated by macroalgae and sessile benthic invertebrates, and are broadly similar to areas of comparable topography to the north (Inner Gneerings offshore from Mooloolaba and offshore of Moreton Bay) and to the south, for example at Julian Rocks offshore of Byron Bay (Fisheries Research Consultants 1991; Harriott et al. 1999; Edwards & Smith 2005; Baronio & Butcher 2008; Fellegara 2008; Schlacher-Hoenlinger et al. 2009).

The fish fauna of the Gold Coast region is similar to that recorded at Flat and Shag Rocks offshore of Moreton Bay, Julian Rocks and the Solitary Islands, offshore of Coffs Harbour and to a lesser extent to the ex-HMAS Brisbane near Mooloolaba (Robinson & Pollard 1982; Parker 1995; Parker 1999; Edwards & Smith 2005; Malcolm et al. 2009; Schlacher-Hoenlinger et al. 2009). The smaller inshore reefs of the Gold Coast region typically support a lower abundance, richness and diversity of reef fish (Edwards & Smith 2005; frc environmental 2005).

Kirra Reef

In April 2014, Kirra Reef covered an area of 2920 m^2 ; predominantly a single area of the northern section of the original reef. The rocky outcrops rose between 1 and 2 m above the clean mobile sand. Several outcrops were found to extend to more than 2 m above the seafloor. There had been a slight reduction in the size of the reef since July 2012, when the reef covered an area of 3200 m^2 .

The benthic assemblages of Kirra Reef have previously been characterised by a high cover of macroalgae and a moderate cover of sessile benthic invertebrates, including a few hard corals (Edwards & Smith 2005; frc environmental 2005). Macroalgae covers the majority of the reef substrate with crinoids (feather stars), ascidians (sea squirts), and sponges, typically the most abundant benthic fauna, whilst anemones, soft corals and urchins are present in low numbers (Fisheries Research Consultants 1995b; a; 1996; frc environmental 2003; 2004; Edwards & Smith 2005; frc environmental 2005). The composition of benthic assemblages at Kirra Reef was broadly similar to that described from adjacent rocky reefs, including Palm Beach Reef (Hollingsworth 1975; Edwards & Smith 2005), and also those of the southern QLD and northern NSW bioregions (refer

Harriott et al. 1999; Baronio & Butcher 2008; Fellegara 2008; Schlacher-Hoenlinger et al. 2009).

Changes in the height of sand around the base of rocky outcrops appears to be a major factor influencing the cover of benthic flora and fauna, periodically resulting in a bare zone on rocks within 0.8 to 1 m of the seafloor. Exposure to wave driven sand scouring and smothering is an important factor influencing the distribution and abundance of sessile species (Kay & Keough 1981; McGuinness 1987). Outcrops on the eastern side of the reef complex, where wave action and sand abrasion are greatest, have historically supported a lower abundance of benthic fauna than outcrops on the northwest side (Fisheries Research Consultants 1995a; b; 1996; frc environmental 2003; 2004; 2005; 2010).

Strong wave action results in sustained abrasion of the dominant brown macroalgae (*Sargassum flavicans* & *Ecklonia radiata*), which causes the fronds to break apart. The continual re-suspension of algal fragments (commonly referred to as 'cornflakes') can dramatically reduce water clarity and visibility. Similar to the July 2012 survey, algal fragments were largely absent in April 2014, which could be due to long periods of relatively benign physical conditions.

Palm Beach Reef

Palm Beach Reef is an extensive rocky reef, lying between the mouths of Tallebudgera Creek to the north and Currumbin Creek to the south. The inner section of the reef is approximately 400 metres off the beach, and lies in 9 to 12 metres of water. Palm Beach Reef lies in slightly deeper water than Kirra Reef, is much larger and has greater topographical relief.

Sessile invertebrates, including sponges, corals and ascidians, dominate the benthic assemblage of Palm Beach Reef (Edwards & Smith 2005; frc environmental 2005; Reef Check 2010). The cover of ascidians, sponges and other invertebrates at Palm Beach Reef has historically been similar to that recorded from the outer sections of Kirra Reef. However, the cover of macroalgae has consistently been lower on Palm Beach Reef than on Kirra Reef. The proximity of Palm Beach Reef to two creek mouths, and the absence of strong currents in the area, typically results in a high level of turbidity. High turbidity together with greater depth and a high abundance of grazing species, such as urchins, is likely to have contributed to a lower cover of macroalgae on Palm Beach Reef.

2 Methods

The methods used in the July 2014 survey were developed for the Stage I survey completed in 1995, and used for subsequent surveys in the 1996, 2001, 2003, 2004, 2005, 2010 and 2012 (following the requirements of the ToR). Data was collected from Kirra Reef and the comparative reef, Palm Beach Reef, further north (Figure 2.1).

The aim of this survey was to assess the effect of the sand bypass project (and subsequent increases in sand load) on the nature and magnitude of change in the cover of floral and faunal, assemblage structure, and the abundance of selected invertebrates.




Figure 2.1 Location of (a) Palm Beach Reef and (b) Kirra Reef

2.1 Benthic Flora and Macroinvertebrates

The number of sites surveyed has varied over time as a result of the fluctuating level of sand surrounding Kirra Reef. A historical recount of sites surveyed can be found in the frc environmental 2010 report *Tweed River Entrance Sand Bypassing Project (TRESBP):* Kirra Reef Ecological Monitoring 2010.

In April 2014, large emergent outcrops were still found only in the northern section of Kirra Reef. In the eastern section there was a small emergent, macroalgal-covered outcrop (2 m²) surrounded by sand in approximately 2 m of water (Figure 2.2). Therefore, it was not possible to compare the eastern sections to the northern sections. The majority of the

reef was largely exposed to a similar level of wave action and sediment deposition and was within a depth range of 5 to 8 m.

Only three sites (KRN1, KRN2, KRO3) were surveyed in April 2014, due to the extent of available reef habitat compared with previous surveys (3,700 m²) as the remainder of the sites previously surveyed were covered by sand (Figure 2.3 and Map 1). The area of reef was determined using ESRI ArcGIS based on a rectified satellite image taken in April 2014 (Near maps 2014). Three comparative sites (PB1, PB2 and PB3) were also surveyed at Palm Beach Reef in a similar depth of water (9 to 12 m) and similar spatial arrangement (Map 3).

Figure 2.2

Small exposed outcrop in the eastern section of Kirra Reef.




Figure 2.3

Diver surveying benthic assemblages at Kirra Reef in April 2014.



At each site, benthic assemblages were surveyed in fifteen 0.25 m² quadrats, and the percent cover of benthic macroalgae, turf algae, sponges, ascidians, hard corals and soft corals were assessed visually. The number of large ascidians (*Pyura stolonifera*), crinoids (feather stars), barnacles, urchins, tubeworms, polychaetes, hydroids, zoanthids and cowries was also noted. These are the same taxonomic groups recorded in the 1995, 1996, 2001, 2003, 2004, 2005, 2010 and 2012 surveys. The dominant species of macroalgae were also recorded, and notes were made on the apparent health of each taxonomic group.

Map 1 Sites surveyed at Kirra Reef in 2014.



Map 2 Site surveyed at Palm Beach in 2014.



2.2 Fish

The species richness and relative abundance of fish at Kirra Reef and Palm Beach Reef were assessed using a combination of underwater visual census (UVC) and video surveys. The combination of these techniques represents the most cost-effective and efficient means of obtaining data on the structure of fish assemblages in different habitats (Murphy & Jenkins 2010). These approaches were the same as used in the 1995, 1996, 2001, 2003, 2004, 2005, 2010 and 2012 surveys. Continuity in the make up of the dive team over the years of monitoring has ensured a high degree of accuracy and precision in fish identification, and in estimating their relative abundance.

2.3 Data Analysis

Due to the limited number of sites sampled, the comparison of the composition of benthic assemblages (cover of each benthic group such as macroalgae, turf algae, corals, sponges and ascidians) over time was restricted to sites KRN1, KRN2 and KRO3 on Kirra Reef and the three comparative sites at Palm Beach Reef (PB1, PB2 and PB3). The analyses completed in 2014 incorporated the data for these six sites from all previous surveys.

Permutational multivariate analysis of variance (PERMANOVA) was used to determine differences in the composition (cover of benthic fauna and taxonomic group) of benthic assemblages between Kirra Reef and Palm Beach Reef over time. This was a new analysis undertaken for the first time in 2012. PERMANOVA is analogous to multivariate analysis of variance (MANOVA); however, rather than using F-tables to derive statistical significance, PERMANOVA uses permutational methods, which require fewer assumptions to be met (Anderson 2001; Anderson et al. 2008). This analysis enables an examination of changes in the community as a whole, which can be more informative than looking at individual components in isolation. In this case, PERMANOVA was used to examine changes in the comparison of benthic assemblages over time.

A three factor PERMANOVA was used to examine differences in the composition of benthic assemblages, with survey (fixed factor), locations (Palm Beach Reef and Kirra Reef, fixed factor) and sites (nested in locations and a random factor) as the factors. Square root transformed data was converted to a Bray Curtis distance matrix and tested for significance using 9999 permutations, where possible. Non-metric multidimensional scaling (nMDS) ordinations were used to visually represent the variation in the composition of assemblages between reefs, separately for each survey.

Analysis of similarity (ANOSIM) was used to determine the magnitude of difference in community composition between the two reefs in each survey. ANOSIM was also used to

determine the difference in community composition between Kirra Reef during the baseline survey in April 1995 and the current survey. ANOSIM is analogous to analysis of variance (ANOVA) in univariate statistics (Clarke 1993; Smith 2003). A global R statistic is calculated to determine whether there is a significant difference among all samples. Comparison of pair-wise R values can give an indication of how different communities are: R values close to 0 indicate little difference, values around 0.5 indicate some overlap and values close to 1 to indicate many or substantial differences. Similarity percentages (SIMPER) analysis determined the key groups that contributed to the differences between surveys.

PERMANOVA can also be used to examine single variables, analogous to ANOVA, except it does not have the same restrictive assumptions that ANOVA has (Anderson et al. 2008). This is because unlike ANOVA, PERMANOVA uses a permutation method to assess significance, which has fewer data assumptions than ANOVA and allows a more accurate assessment of unbalanced experimental designs or where biological data is not normally distributed. Therefore in 2014, separate univariate PERMANOVAs (rather than ANOVAs) were used to compare differences in the cover of macroalgae, turf algae and the abundance of crinoids and ascidians. These univariate analyses were completed using the design described above (Anderson et al. 2008).

Further information on the use and interpretation of PERMANOVA and other analyses used in this report is provided in Appendix A.

3 Results

3.1 Cover of Benthic Assemblages

The composition of benthic fauna and flora (% cover and type combined) at Kirra Reef and Palm Beach Reef changed between surveys, with the magnitude of difference between reefs varying over time (Table 3.1 to Table 3.3). There were substantial differences in the composition of the benthic assemblages between the two reefs in each survey from 1995 to 2014 (Figure 3.1 to Figure 3.3); however, benthic assemblages at the two reefs were most different from each other in 1996, 2001 and 2005 (note the larger R statistic values in Table 3.2; and the separation of reefs in Figure 3.1 & Figure 3.2). In 2010 and 2012, there was greater overlap in the composition of assemblages between reefs (Table 3.2 and Figure 3.1 & Figure 3.2). In 2014, assemblages on each reef were distinct from each other but there was some overlap, similar to the results in 1995 (Table 3.2, Figure 3.1 & Figure 3.3). In 2014, the main difference in the composition of benthic assemblages between Kirra and Palm Beach reefs was due to a lower cover of soft corals, hard corals and ascidians at Kirra Reef (Table 3.3).

Table 3.1 PERMANOVA results for multivariate differences in the composition of benthic assemblages between surveys and locations.

Factor	df	MS effect	Pseudo-F	p (perm)
location	1	3.12 x 10 ⁵	133.42	0.001
survey	8	36506	9.62	0.001
site (location)	4	2365	3.1	0.001
location x survey	8	17540	4.62	0.001
site (location) x survey	30	3810	5	0.001
error	818	763		

Shading denotes significance at p < 0.05

Table 3.2 Pairwise ANOSIM results for differences in assemblage composition between reefs for each survey.

Survey	1995	1996	2001	2003	2004	2005	2010	2012	2014
ANOSIM R statistic	0.577	0.713	0.797	0.661	0.543	0.932	0.509	0.484	0.594
p value	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001

Shading denotes significance at p < 0.05

Table 3.3 PERMANOVA post hoc pairwise results for the differences between locations in each survey events, for the assemblage composition and cover of selected benthic groups.

Year		mblage oosition	Macr	oalgae	Turf	Algae	Soft	Corals	Hard	Corals	Sponges		Asci	Ascidians	
	t	p (MC) ^a	t	p (MC) ^a	t	p (MC) ^a	t	p (MC) ^a	t	p (MC) ^a	t	p (MC) ^a	t	p (MC) ^a	
1995	6.05	0.001	0.34	0.851	1.50	0.179	3.67	0.015	3.47	0.033	9.64	0.001	5.29	0.005	
1996	3.58	0.001	0.53	0.668	4.81	0.007	1.81	0.132	4.57	0.005	0.73	0.553	2.38	0.072	
2001	3.80	0.004	6.84	0.001	3.79	0.008	1	0.388	_	_	3.90	0.013	2.35	0.070	
2003	2.99	0.019	7.06	0.004	6.28	0.001	6.73	0.002	1.73	0.153	0.09	0.974	2.53	0.068	
2004	4.62	0.002	1.04	0.368	2.99	0.014	0.59	0.618	7.38	0.001	0.75	0.531	1.75	0.159	
2005	7.02	0.001	0.64	0.635	10.25	0.002	2.01	0.118	1.49	0.191	4.24	0.008	0.76	0.530	
2010	3.60	0.001	0.94	0.391	0.60	0.626	6.73	0.001	2.48	0.064	6.11	0.001	0.61	0.612	
2012	3.04	0.007	2.61	0.045	0.93	0.438	10.41	0.001	10.59	0.001	2.68	0.050	5.96	0.002	
2014	3.14	0.002	0.49	0.682	0.74	0.528	3.59	0.014	9.42	0.001	0.36	0.798	2.74	0.025	

Shading denotes significance at p < 0.05

a p values based on Monte Carlo tests




Figure 3.1 Multi-dimensional scaling plot of benthic cover in the 1995, 1996, 2001 and 2003 surveys.




Figure 3.2 Multi-dimensional scale plot of benthic cover in the 2004, 2005, 2010 and 2012 surveys.




Figure 3.3 Multi-dimensional scale plot of benthic cover in the 2014 survey.

The benthic assemblages on Kirra Reef were significantly different between the baseline survey in April 1995 and the survey in April 2014 (ANOSIM, Global R = 0.851, p = 0.001, Figure 3.4). Differences in benthic assemblages between surveys were due to a:

- · lower cover of bare substrate in 2014
- · higher cover of ascidians in 2014, and
- · higher cover of turf algae in 2014 (SIMPER analysis).

The benthic assemblages recorded in April 2014 were representative of a community in succession. There was no cover of ascidians or crustose coralline algae recorded at Kirra Reef in April 1995, but they were present in April 2014. Similarly hard and soft corals were present in April 1995, but hard corals were not recorded in April 2014, and soft corals were rare.




Figure 3.4 Multi-dimensional scale plot of benthic cover at Kirra Reef in April 1995 and April 2014.

3.2 Benthic Algae

Macroalgae

In April 2014, as in previous surveys, the macroalgae *Sargassum* sp. dominated the benthic assemblages at all sites at Kirra Reef (Figure 3.5). Other species present included:

- Dictyopteris arostichoides
- Dilophus intermedius
- Zonaria sp.
- · Laurencia brongniartic
- Amphiroa anceps
- · Caulerpa lentillifera, and
- · Halimeda discoidea.

The mean cover of macroalgae at Kirra Reef has significantly declined over time, with the greatest magnitude of decline recorded between January 2001 and May 2003 (Table 3.4, site (location) x survey interaction). In surveys prior to May 2003, *Sargassum* sp. formed dense carpets over the rocky substrate, covering up to 58% of the available surface area on the reef. There was a distinct decline in the cover of macroalgae between January 2001 and May 2003, which appears strongly correlated with the decrease in reef area during that time as well as a decline in the cover of *Sargassum* sp.. Overall, *Sargassum* sp. dominated macroalgal communities in each survey, except in 2003 and 2004, and this species may be an important indicator species on Kirra Reef. Since February 2010, the cover of all macroalgae has increased from the 12% recorded in February 2010 to 25% in April 2014, which is similar to the cover recorded in April 1995 (Figure 3.6). In contrast, the cover of macroalgae on Palm Beach Reef was consistently lower than Kirra Reef, generally being less than 5% of the available surface area (Figure 3.6).

The macroalgae species recorded at Palm Beach Reef, included:

- Amphiroa anceps
- Laurencia brongniartic
- Chlorodesmis major, and
- Zonaria sp..

Figure 3.5

Sargassum sp. dominated the macroalgal communities of Kirra Reef at site KN3 in April 2014.






Figure 3.6 Mean cover of macroalgae (\pm SE) at Kirra Reef and Palm Beach Reef in all surveys.

Table 3.4 Univariate PERMANOVA results for differences in the cover of macroalgae between surveys and location.

Factor	df	MS effect	Pseudo-F	p (perm)
location	1	15087	2.52	0.079
survey	8	15019	1.48	0.174
site (location)	4	5995	12.68	0.001
location x survey	8	22158	2.19	0.056
site (location) x survey	32	10128	21.41	0.001
error	756	473		

Shading denotes significance at p < 0.05

Turf Algae

The mean cover of turf algae at both Kirra Reef and Palm Beach Reef has varied considerably between surveys, between reefs and between sites (Table 3.5). Both Kirra and Palm Beach reefs showed major increases in the percent cover of turf algae from 2010 to 2012, but then a decrease in cover between 2012 and 2014 (Figure 3.8). The cover of turf algae is typically lower at Kirra Reef than at Palm Beach Reef. However, in April 2014, turf algae cover was similar between the two reefs (Table 3.3).

Increased cover of turf algae is typically related to good light conditions, high concentrations of nutrients and low numbers of grazing species such as fish and sea urchins, and perhaps reflects a more physically robust growth form suited to high wave energy environments (than foliose macro-algae).

Figure 3.7

Areas of Palm Beach Reef support little macroalgae but have a high cover of turf algae (April 2014).






Figure 3.8 Mean cover of turf algae (\pm SE) at Kirra Reef and Palm Beach Reef in all surveys.

Table 3.5 Univariate PERMANOVA results for differences in the cover of turf algae between surveys and locations.

Factor	df	MS effect	Pseudo-F	p (perm)
location	1	83788	23.07	0.113
survey	8	13335	4.98	0.001
site (location)	4	3632	8.76	0.001
location x survey	8	14479	5.40	0.001
site (location) x survey	32	2680	6.46	0.001
error	756	415		

Shading denotes significance at p < 0.05


3.3 Benthic Macro-Invertebrates

Sponges

The mean cover of sponges (Figure 3.9) at Kirra Reef declined significantly from more than 20% in March 2004 to less than 1% in February 2010. Since 2010, the cover of sponges has increased slightly, with sponges covering 4% of the reef in April 2014 (Figure 3.10). In April 2014, the cover of sponges was higher than recorded during the baseline survey in April 1995. The mean cover of sponges at Palm Beach Reef was significantly higher than at Kirra Reef, except in April 1995, June 1995, January 2001, February 2005 and February 2010 (Figure 3.10 and Table 3.3).

Figure 3.9

Sponges were much more abundant at Palm Beach Reef than at Kirra Reef since 2005.






Figure 3.10 Mean cover of sponges (\pm SE) at Kirra Reef and Palm Beach Reef on all surveys.

Table 3.6 Univariate PERMANOVA results for differences in the cover of sponges between surveys and locations.

Factor	df	MS effect	Pseudo-F	p (perm)
location	1	82369	30.64	0.118
survey	8	8867	3.12	0.008
site (location)	4	2688	3.68	0.003
location x survey	8	12972	4.56	0.001
site (location) x survey	32	2844	3.89	0.001
error	756	731		

Ascidians

In April 2014, the mean abundance (individual per $0.25 \, \text{m}^2 \pm \text{SE}$) of the ascidian, *Pyura stolonifera* (Figure 3.11), on Kirra Reef was higher than at Palm Beach Reef (Figure 3.12 and Figure 3.12). The slight increase in mean abundance of *P. stolonifera* at Kirra Reef in April 2014 is due to increased recruitment and survival in the years prior to those surveys.

The cover of all ascidians other than *P. stolonifera* has varied considerably over time (Figure 3.13 and Table 3.7). In April 2014, the mean cover of all other ascidians on Kirra Reef and Palm Beach Reef was higher than previously reported in the baseline survey in April 1995 (when no ascidians were recorded).

Figure 3.11

Ascidians (*Pyura stolonifera* and *Cnemidocarpa stolonifera*) were present at Kirra Reef in April 2014.






Figure 3.12 Mean abundance of ascidians ($Pyura\ stolonifera$) (individuals / 0.25 m²) (\pm SE) at Kirra Reef and Palm Beach Reef in all surveys.




Figure 3.13 Mean cover of ascidians (\pm SE), other than *Pyura stolonifera*, at Kirra Reef and Palm Beach Reef in all surveys.

Table 3.7 Univariate PERMANOVA results for the differences in cover of all ascidians other than *Pyura stolonifera* between surveys and locations.

Factor	df	MS effect	Pseudo-F	p (perm)
location	1	1998	1.57	0.381
survey	8	22264	8.73	0.001
site (location)	4	1276	2.09	0.074
location x survey	8	17109	6.71	0.001
site (location) x survey	32	2551	4.18	0.001
error	756	611		

Hard Coral

The cover of hard coral is generally low on Kirra Reef, being less than 2% of the available surface area. In April 2014, this trend continued with no hard corals identified during the survey on Kirra Reef (Figure 3.14). Hard coral often covered more of the available surface area on Palm Beach Reef than at Kirra Reef over time, except during the baseline survey in April 1995, when the cover of hard coral was low on both reefs (Figure 3.14 and Table 3.8, PERMANOVA location x survey interaction).




Figure 3.14 Mean cover of hard corals (\pm SE) at Kirra Reef and Palm Beach Reef in all surveys.

Table 3.8 Univariate PERMANOVA results for the differences in the cover of hard coral between surveys and locations.

Factor	df	MS effect	Pseudo-F	p (perm)
location	1	76173	305.49	0.111
survey	8	8480	15.87	0.001
site (location)	4	249	0.62	0.681
location x survey	8	6654	12.45	0.001
site (location) x survey	32	534	1.34	0.109
error	756	399		

Soft Coral

In April 2014, the mean cover of soft coral at Kirra Reef was less than 1%. Although the cover was low, this was the first time that soft corals had been recorded at Kirra Reef since 2005 (Figure 3.15). Prior to being covered with sand, Kirra Reef up to 10% cover of soft corals. Soft corals have historically covered more of the available space on Palm Beach Reef than on Kirra Reef. The cover of soft coral at Palm Beach Reef has varied substantially over time, being highest in April 1995 (48%) and has significantly declined since that time; however, the cover of soft coral has been increasing since February 2005 (Figure 3.15).




Figure 3.15 Mean cover of soft corals (\pm SE) at Kirra Reef and Palm Beach Reef in all surveys.

Table 3.9 Univariate PERMANOVA results for the differences in the cover of soft coral between surveys and locations.

Factor	df	MS effect	Pseudo-F	p (perm)
location	1	110360	82.84	0.096
survey	8	11370	8.32	0.001
site (location)	4	1332	2.44	0.041
location x survey	8	6051	4.43	0.005
site (location) x survey	32	1367	2.51	0.001
error	756	545		

Crinoids

The mean abundance (individuals / 0.25 m²) of crinoids at Kirra Reef and Palm Beach Reef has declined considerably since the baseline survey in 1995 (Table 3.10 & Figure 3.17). However, the mean abundance of crinoids on Kirra Reef has been increasing since February 2005 (Figure 3.17). In April 2014, the abundance was extremely patchy over Kirra Reef, with most individuals found in crevices or under ledges (Figure 3.16). In comparison, no crinoids were recorded at Palm Beach Reef. Crinoids are often found on new reef structures in the region, where they quickly recruit and cling to bare hard surfaces (Schlacher-Hoenlinger et al. 2006).

Figure 3.16

Crinoids (feather stars) were patchily distributed found at Kirra and Palm Beach reefs in April 2014.






Figure 3.17 Mean abundance of crinoids (individuals / $0.25~\text{m}^2$) (\pm SE) at Kirra Reef and Palm Beach Reef in all surveys.

Table 3.10 Univariate PERMANOVA results for the differences in crinoid density between surveys and locations.

Factor	df	MS effect	Pseudo-F	p (perm)
location	1	5947	28.81	0.09
survey	8	1340	2.42	0.036
site (location)	4	206	1.16	0.308
location x survey	8	3890	7.03	0.001
site (location) x survey	32	554	3.11	0.001
error	756	178		

3.4 Fish

The species richness of fish recorded from Kirra Reef across all monitoring events has ranged from 12 to 53. In April 2014, 53 species were identified at the reef, which was the highest recorded in any survey (Figure 3.18). Species richness at Palm Beach Reef was slightly lower (39 species) and the variation between sampling periods was generally less than at Kirra Reef. Nonetheless, the fish assemblages as a whole were quite similar between both reefs.

In April 2014, the assemblage of fish at Kirra Reef comprised species from all trophic levels, including detritivores, planktivores, herbivores and carnivores (Appendix B). As in previous surveys, the assemblage was dominated by herbivores and planktivores. Yellowtail (*Trachurus novaezelandiae*), Australian mado (*Atypichthys strigatus*), stripeys (*Microcanthus strigatus*), sweep (*Scorpis lineolatus*) and various wrasses were very abundant and remained the dominant species (Figure 3.19 and Appendix B). One new species was recorded at both Kirra Reef and Palm Beach Reef that was not recorded in any previous survey, the anthias (*Anthias* sp.). Two new species were recorded at Palm Beach Reef, the Queensland grubfish (*Parapercis queenslandiae*) and barred soapfish (*Diploprion bifasciatum*) (Figure 3.20). An additional three new species were recorded at Kirra Reef:

- · barracuda (Sphyraena argentea)
- cowtail stingray (Pastinachus atrus) (Figure 3.21), and
- · Moorish idol (Zanclus cornutus).

Moon wrasse (*Thalassoma lunare*), neon damsels (*Pomacentrus coelestis*), large-scaled parma (*Parma oligolepis*) (Figure 3.22), Australian mado (*Atypichthys strigatus*), and yellowtail (*Trachurus novaezelandiae*) continued to be abundant at Palm Beach in April 2014. The complete list of species recorded and the relative abundance is presented in Appendix B.




Figure 3.18 Number of fish species recorded at Kirra Reef and Palm Beach Reef on each survey.

Figure 3.19

Yellowtail were abundant at Kirra
Reef in April 2014.



Figure 3.20

Barred soapfish were recorded at Palm Beach Reef for the first time in April 2014.



Figure 3.21

Cowtail stingray was recorded at Kirra Reef for the first time in April 2014.



Figure 3.22

Large scale parma continued to be abundant at Palm Beach Reef in April 2014.



4 Discussion

4.1 Changes to the Ecological Condition of Kirra Reef

The greatest change to the ecological condition of Kirra Reef has been the loss of large areas of hard substrate that support benthic flora and fauna, due to burial by shifting sand. The delivery of large sand volumes during the Initial TRESBP operations resulted in a significant increase to the beach width at Kirra, with wave action and tidal currents redistributing some of this sand over Kirra Reef.

Between August 2008 and August 2009, there had been a loss of more than 200,000 m³ of sand from the upper beach and offshore area between the Kirra Point groyne and the Miles Street groyne. About 60,000 m³ of this sand was lost from the beach above the water line, and about 160,000 m³ was lost in shallow waters out to about 6 metres depth. By mid-2009, seabed levels in the nearshore area had reduced by up to about 2 metres and there was an increase in the extent of reef habitat, with a large increase in the area of northern section of rocky reef occurring between February 2010 and July 2012.

In April 2014, the rocky outcrops in the northern section of Kirra Reef supported a moderately diverse benthic assemblage, dominated by macroalgae, and high fish diversity. The ecological assemblage exhibited signs of ongoing stress from physical disturbance, including a reduced cover of hard coral and soft coral, and greater variation in the cover of sponges relative to both previous surveys and the cover of similar organisms at Palm Beach Reef. However, Kirra Reef continues to provide habitat to a range of flora and fauna, and provides important marine ecological functions and services in the region. It is possible that as sand levels have stabilised over the past two years, assemblages are beginning to become more similar to those recorded prior to implementation of the TRESBP. For example, there was a decrease in the cover of turf algae and an increase in the cover of macroalgae from 2012 to 2014; an increase in the cover of sponges, ascidians and crinoids from 2012 to 2014; and soft corals were recorded at Kirra Reef in 2014 for the first time since 2005. However, assemblage composition is still significantly different to that recorded in the baseline surveys.

Benthic Macroalgae

The cover of macroalgae at Kirra Reef increased slightly between the July 2012 and April 2014 surveys; however, it remained well below the peak of 60% cover recorded in January 2001. The cover of macroalgae declined at Palm Beach Reef over the same period. However, the cover of macroalgae at Palm Beach Reef was generally much lower

than at Kirra Reef over time. There are several possible reasons for the lower cover of macroalgae at Palm Beach including:

- Palm Beach Reef is slightly deeper than Kirra Reef and generally has higher turbidity, and therefore greater light attenuation. The quantity and quality of available light affects the distribution and growth of macroalgae (Miller & Etter 2008).
- Differences in the pattern of recruitment of algal species to the reefs due to different water currents and timing of the surveys (Kennelly 1987b).
- Increased competition with turf algae and sessile benthic invertebrates, which compete for space with macroalgae (Kennelly 1987a; Miller & Etter 2008).
- Presence of different species and higher density of herbivores at Palm Beach Reef (particularly sea urchins and herbivorous fish), which would graze on macroalgae (McCook 1997; Jompa & McCook 2002).

Temporal variation in the cover of macroalgae at Kirra Reef is likely to be principally due to the effect of physical disturbance from wave action. The negative effect of physical disturbance on the cover of macroalgae is evident during survey events when algal fragments form a dense covering over the reef. Increased smothering by sand can reduce the diversity (Hatcher et al. 1989), abundance, recruitment, growth, survival and seasonal regeneration of macroalgae (Umar & Price 1998; Cheshire et al. 1999). Biological assemblages exposed to physical stress also typically exhibit greater levels of temporal and spatial variability (Warwick & Clarke 1993; Chapman et al. 1995).

An alternative explanation for the changes in the cover of macroalgae on Kirra Reef over time is that as the extent of the reef has changed; the fish associated with the reef became more concentrated, increasing the grazing pressure. Increased grazing pressure from fish and mobile invertebrates can reduce the coverage and diversity of macroalgae on reefs (McCook 1997; Jompa & McCook 2002).

The cover of turf algae at both Kirra and Palm Beach reefs also varied significantly between surveys, typically covering less of the available surface area on Kirra Reef than Palm Beach Reef. However, in April 2014 the cover of turf algae was similar between reefs and lower than recorded in 2012. The lower cover of turf algae at Kirra Reef was likely due to the increased cover of macroalgae, sponges and ascidians, while at Palm Beach Reef this was likely due to an increase in hard coral cover. However, the relationship between algal dynamics, physical disturbance, water quality and herbivore grazing activity is complex, and the cover of turf algae can exhibit extreme temporal variability as a consequence of the interaction between top-down and bottom-up processes (Russ 2003; Bellwood et al. 2006; Hughes et al. 2007; Albert et al. 2008; Hoey

& Bellwood 2008; Mumby 2009). Further investigation would be required to determine the exact mechanisms of change in both macroalgae and turf algae assemblages on these reefs.

Benthic Macroinvertebrates

The cover of benthic macroinvertebrates varied between Kirra Reef and Palm Beach Reef. The cover of sponges, hard coral and soft coral was higher at Palm Beach Reef, while the cover of ascidians and density of crinoids was higher at Kirra Reef. At Kirra Reef, the continued burial and re-emergence of rocky outcrops (mediated by the sand bypassing project) is likely to have increased temporal variability in the distribution and abundance of benthic macroinvertebrates. Additional perturbations such as wave action and sand abrasion are likely to have resulted in the decline in the cover and diversity of the benthic macroinvertebrates at Kirra Reef between some surveys particularly between March 2004 and February 2005. Benthic macroinvertebrates such as ascidians, sponges, hard coral and soft coral, are highly susceptible to the effects of storm and wave disturbance, physical abrasion and burial by sand (Kay & Keough 1981; Walker et al. 2008), which affects settlement, growth rates and survival of these taxa (Dodge & Vaisnys 1977; Rogers 1990).

Physical disturbance from sand burial, sand abrasion and the action of storm waves appear to keep the benthic assemblages on Kirra Reef in a state of early succession. It is common for early pioneer species, such as macroalgae or barnacles, to recruit rapidly to a hard surface in large numbers, allowing these species to dominate assemblages early in the successional trajectory (Walker et al. 2007). Another indicator of the early state of succession at Kirra Reef is that the cover of hard and soft coral has remained very low. In April 2014, soft coral was recorded at Kirra Reef for the first time since February 2005. There was no hard coral recorded on Kirra Reef, which was likely related to increased physical disturbance from sand burial and abrasion and the loss of some reef area since 2012, as well as competition and recruitment processes as discussed below.

Benthic macroinvertebrate cover may be affected indirectly through increased competition with macroalgae for space. The presence of large macroalgae can affect the recruitment and survival of sessile benthic invertebrates as fronds moving with wave action, sweep and abrade the surface of rocks, killing new recruits, especially corals (Kennelly 1989; McCook et al. 2001). It can take several years for hard and soft coral to become dominant on reefs in the region (Schlacher-Hoenlinger et al. 2009). Therefore, we would not expect hard coral to become abundant until several years after the reef had been uncovered, and only if the physical disturbance regime and supply of new recruits was sufficient to support survival of these species. It should be noted that the cover of hard

and soft corals at Kirra Reef was both very low and very patchy even prior to the start of the sand bypassing operations.

Due to increased wave action and sedimentation mediated by the sand bypassing project, the mean cover of ascidians and sponges was expected to be much lower on Kirra Reef than Palm Beach Reef. This pattern was observed in the cover of sponges in 2014, which covered 4% of Kirra Reef and 7% on Palm Beach Reef, although the magnitude of difference in cover was smaller than in 2012. However, this pattern was not consistent with ascidians (other than *P. stolonifera*), which covered 20% of Kirra Reef, compared with 9% cover on Palm Beach Reef. The increasing cover of sponges and ascidians on Kirra Reef between 2012 and 2014 indicates that Kirra Reef is slowly recovering from the increased stress from physical disturbance such as increased wave action and physical abrasion from sand scour. Sponges and ascidians are highly susceptible to smothering and sand abrasion, unless they have a thick tunic (outer covering made of keratin) like the ascidian *P. stolonifera*, or strong internal keratin, silicon or calcareous structures in the case of some species of sponges (Kay & Keough 1981; McGuinness 1987; Walker et al. 2008).

Fish

The reduction in area of hard substrate at Kirra Reef over time has affected the composition of fish assemblages; however, the reef continues to support a high diversity of reef-associated, and pelagic (i.e. non-reef associated) fish species. This indicates that despite its diminished size from 1995 conditions, Kirra Reef continues to provide valuable habitat for a variety of fish species from different functional groups.

There is a high degree of inter-annual variability in the species and abundance of the fish present at Kirra Reef compared with the assemblage at Palm Beach Reef, which is likely to reflect temporal variability in the available habitat as a consequence of reef burial and re-emergence. The diversity, quality and areal extent of reef habitat are the most important factors influencing the distribution, abundance, biomass and diversity of reef fish (Bellwood & Hughes 2001; Friedlander et al. 2003). Diversity and abundance of fish can increase with greater structural complexity and increased heterogeneity of available habitats (Bellwood & Hughes 2001). Logically, this suggests that periods of chronic reef burial may reduce the overall diversity of reef-associated fish species. Despite the reduction in overall reef space, there was a larger variety of habitats in 2014, which was observed to have a positive effect on the diversity of fish assemblages. This is particularly important for several species that depend on the presence of reef habitat (i.e. oldwife, moray eels, damselfish and Australian mado that were recorded in April 2014, but were not recorded in July 2012).

The abundance and diversity of fish are likely to be lowest following periods of severe weather, which create unfavourable conditions for many species, and may further exacerbate the affects of sedimentation. The biomass of fish is known to decrease with increasing exposure to physical disturbance from wave action and strong currents (Friedlander et al. 2003). As the surveys were completed at different times of the year, variation in the prevailing conditions at the time of sampling could also influence the types of fish observed and the amount of reef habitat that is available at any time. Many of the species may also be affected by seasonal changes in the water temperature, such as damselfish, which are less abundant in cool waters. However, the overall diversity of the fish assemblage at Kirra Reef is likely to reflect the complex interactions between physical disturbance, food availability and competition, local weather and sea conditions.

4.2 Impacts of the Sand Bypassing System on Kirra Reef

In addition to assessing changes to the ecological condition of Kirra Reef over time, frc environmental was commissioned to assess any noticeable impacts of the TRESBP on the extent of the reef, taking into account the impacts predicted in the:

- Tweed River Entrance Sand Bypassing Project Permanent Bypassing System Environmental Impact Statement / Impact Assessment Study, prepared by Hyder Consulting, Paterson Britton & Partners Pty Ltd and WBM Oceanics Australia Joint Venture in June 1997
- Impact Assessment Review Report for Tweed River Entrance Sand Bypassing Project Permanent Bypassing System, prepared by the Queensland Department of Environment in March 1998, and
- Report on Historic Changes at Kirra Beach, prepared by P.K. Boswood and R. J. Murray of the then Queensland Department of Environment in March 1997.

The EIS / IAS is a document of broad scope. Our comparison of predicted and actual impacts on Kirra Reef has focused on three principle sections of the EIS / IAS relevant to the consideration of the ecology of Kirra Reef: 'The Existing Environment', 'The Environmental Impacts' and 'Environmental Management'. A more comprehensive review of these sections can be found in our previous monitoring reports (frc environmental 2003; 2004; 2005). In summary, the predicted impacts included accretion of sand around the base of the rock outcrops at Kirra Reef, causing a reduction in extent of the uncovered area of reef. It was predicted that sand delivery as part of the project would eventually mimic 'natural' patterns of sand dispersal, and that the reef would reduce in size to its natural extent (i.e. the extent prior to development of the Tweed River training walls). However, the EIS did not predict the ecological consequences of both the reduced areal

extent or increased wave energy (a consequence of decreased depth) that occurred. Presumably the benthic flora and fauna assemblages of the reef would be expected to return to a condition consistent with the historical reef extent and natural sand transport patterns (and associated coastal fluctuations, wave action, sedimentation and water quality) that were observed in the vicinity of the reef prior to the development of the Tweed River training walls.

The current extent of Kirra Reef, which has been present throughout much of the last decade, is quite different to that of the late 1980s / early 1990s. In April 2014, the areal extent of Kirra Reef was similar to that found in the 2001 survey. When compared with the historical extent of the reef, the current extent of the northern section of the reef most closely approximates that recorded in 1930. However, there has been a consistent decline in the extent of the eastern and inner western sections of the reef, due to increased sand deposited as part of the TRESBP project. Therefore, the extent of Kirra Reef in April 2014 is broadly in accordance with predictions made in the EIS.

While the extent of the reef continues to change over time, the delivery of sand as part of the project now closely matches the natural rate of longshore sand transport, so short-term and seasonal changes in the extent of the reef are now more likely to be the result of wave and current action than a discrete impact of the sand bypassing activity. Ongoing monitoring to assess the physical and ecological dynamics of Kirra Reef will be valuable to determine the magnitude and frequency of changes in the diversity of the benthic assemblages found on Kirra Reef under a mature sand-bypassing regime.

4.3 Impacts of Storms & Seasonality on Kirra Reef

Sessile benthic assemblages on Kirra Reef are highly susceptible to the influence of storms, and associated wave action (Kay & Keough 1981; Walker et al. 2008). The shallow reef is surrounded by mobile sand, which can shift naturally in response to wave action during storms causing burial of large sections of Kirra Reef. This effect has reduced the extent of rocky substrate available for colonisation, and also the availability of refuge habitats, such as crevices and overhangs, which are sheltered from wave action and sand abrasion. In April 2014, there was a slight decrease in reef extent due to a portion of the western edge of the reef being covered by sand.

Wave height typically increases during storm events, and given the shallow depth of Kirra Reef, waves are more likely to shoal and break across the reef during storms. This increases the physical disturbance, abrasion and sedimentation of benthic assemblages on Kirra Reef. Storm disturbance can cause local reductions in the species richness and abundance of coral (Woodley et al. 1981; Massel & Done 1993; Hughes 1994; Connell et al. 1997), and can alter fish assemblages indirectly through habitat modifications

(Kaufman 1983; Jones & Syms 1998) or directly by increasing fish mortality (Lassig 1983). The hydrodynamic forces produced by wave action are an important source of disturbance in intertidal habitats, inflicting damage through direct physical impact and abrasion (Paine & Levin 1981; Denny 1983; Shanks & Wright 1986; Bell & Denny 1994; Walker et al. 2008).

Storm and wave action (and associated sedimentation and abrasion) continue to be important forces shaping the distribution and abundance of benthic species at Kirra Reef. Increased magnitude and frequency of physical disturbance, resulting from increased exposure or susceptibility to storms and associated wave action (as on Kirra Reef), can lead to a decrease in the diversity of sessile invertebrate assemblages (Walker et al. 2008). Disturbance-driven reductions in biodiversity have the potential to impact negatively on the health and productivity of reef ecosystems (Walker et al. 2008). Biodiversity is key to reef health given that many of these species (e.g. sponges, bryozoans and ascidians) contribute a range of vital ecosystem services to reefs, including: nutrient cycling (Scheffers et al. 2004), trophic interactions and food webs (Lesser 2006; Pawlik et al. 2007), bio-erosion (Rutzler 2002; Lopez-Victoria et al. 2006), and stabilizing substrata (Diaz & Rutzler 2001; Wulff 2001).

The impacts of increased wave action and sedimentation on the benthic assemblages at Kirra Reef are likely to be greatest during and immediately following storm conditions. Partitioning the influence of storm and wave driven disturbance, from that of the operation of the TRESBP, would require a much more statistically powerful, and temporarily replicated experimental monitoring design.

4.4 Long-term Impacts of the Sand Bypassing System on Kirra Reef

In April 2014, Kirra Reef covered an area of approximately 40 to 45% (2920 m²) of the reef area present in 1995 (i.e. prior to the operation of the TRESBP). This is largely due to the reduction in the inner western and eastern sections of the reef, which have yet to re-emerge (Figure 1.2). Logically, the loss of reef habitat has dramatically reduced the availability of hard substratum available for colonisation and consequently the diversity of benthic sessile assemblages.

While there is sufficient structure available, we expect that the assemblage of fish will continue to resemble the historical assemblage at Kirra Reef and also that of Palm Beach Reef, perhaps with the inclusion of a greater proportion of cryptic benthic species as they recruit to Kirra Reef. Given that fish are mobile, the greatest effects on these assemblages are likely to be short-term (reversible) changes due to the prevailing conditions and changes to the extent of available habitat to provide shelter and food.

5 Conclusions

The current extent of Kirra Reef is broadly in accordance with predictions made in the EIS. Initial 'catch-up' bypassing of sand resulted in the substantial burial of Kirra Reef. 'Maintenance' bypassing more closely reflects natural patterns of long-shore sand transport and is gradually allowing the reef to re-emerge.

The extent of the reef declined slightly between 2012 and 2014, however apart from some minor burial, the extent of reef exposed has been relatively stable. This has allowed for some succession in the benthic assemblages, resulting in a reduction in turf algae cover and a corresponding increase in the cover of sponges, ascidians, crinoids and the first record of soft corals since 2005. However, while assemblages appear to be moving towards being more similar to the baseline conditions, they are still significantly different to those conditions recorded in April 1995. With the delivery of sand more closely matching the natural rate, it is expected that the reef may undergo short-term changes in extent due to seasonal shifts in sand delivery; however, the diversity of benthic assemblages on Kirra Reef should increase gradually over time, especially if a larger extent of the rocky reef becomes uncovered. This would create more space for a greater suite of species to recruit, and reduces the likelihood of sand abrasion and wave damage. It was expected that the areas of Kirra Reef that were newly exposed in 2012 would undergo a shift from turf-dominated communities to more diverse communities that included macroalgae, sponges, ascidians and potentially hard and soft coral. With the exception of hard corals (which are still absent from Kirra Reef, and are likely to take more time to recolonise), this prediction was confirmed by the results of the April 2014 survey despite the minor burial of hard substrates.

The diversity of fishes associated with Kirra Reef is broadly similar to that recorded prior to the commencement of sand bypassing. Given that fish are mobile, the greatest impacts on fish assemblages are likely to be short-term changes due to the prevailing conditions and changes to the extent of available habitat that provides shelter and food for a variety of different species. If Kirra Reef continues to increase in size there is likely to be a greater proportion of cryptic benthic species as these species are more typical of assemblages in a later stage of succession.

The timeframe for the 'recovery' of communities to this state is currently unknown, and will depend on ambient environmental conditions. Ongoing monitoring will provide insight into the rate of 'recovery' of communities. It is recommended that monitoring be repeated in 2015 (between February and April, when most of the previous surveys have been undertaken) to confirm the results of the April 2014 survey and account for seasonal differences. It is also recommended that surveys be undertaken in 2015 in order to monitor the rate of recovery of Kirra Reef. If communities surveyed in 2015 demonstrate

consistent results, monitoring could be completed bi-annually to monitor succession in the assemblages, unless triggered by a 15% change in the extent of the exposed reef.

6 References

- Albert, S., Udy, J. & Tibbetts, I., 2008, 'Responses of algal communities to gradients in herbivore biomass and water quality in Marovo Lagoon, Solomon Islands', *Coral Reefs* 27: 73-82.
- Anderson, M.J., 2001, 'A new method for non-parametric multivariate analysis of variance', *Austral Ecology* 26: 32-46.
- Anderson, M.J., 2004. PERMDISP: a FORTRAN computer program for permutatinoal analysis of multivariate dispersions (for any two-factor ANOVA design) using permutational tests. Department of Statistics, University of Auckland, New Zealand.
- Anderson, M.J., Gorley, R.N. & Clarke, K.R., 2008. *PERMANOVA+ for PRIMER: Guide to Software and Statistical Methods*. PRIMER-E, Plymouth, UK.
- Baronio, M.A. & Butcher, D.J., 2008, 'Artificial crevice habitats to assess the biodiversity of vagile macro-cryptofauna of subtidal rocky reefs', *Marine and Freshwater Research* 59: 661-670.
- Bell, E.C. & Denny, M.W., 1994, 'Quantifying "wave exposure": a simple device for recording maximum velocity and results of its use at several field sites', *Journal of Experimental Marine Biology & Ecology* 181: 9-29.
- Bellwood, D. & Hughes, T., 2001, 'Regional-scale assembly rules and biodiversity of coral reefs', *Science* 292: 1532-1535.
- Bellwood, D.R., Hughes, T.P. & Hoey, A.S., 2006, 'Sleeping functional group drives coral-reef recovery', *Current Biology* 16: 2434-2439.
- Boswood, P.K. & Murray, R.J., 1997, *Report on historical changes at Kirra Beach*, report prepared for Queensland Department of Environment.
- Cannon, L.R., Goeden, G.B. & Campbell, P., 1987, 'Community patterns revealed by trawling in the inter-reef regions of the Great Barrier Reef', *Memoirs of the Queensland Museum* 25: 45-70.
- Chapman, M.G., Underwood, A.J. & Skilleter, G.A., 1995, 'Variability at different spatial scales between a subtidal assemblage exposed to the discharge of sewage and two control assemblages.', *Journal of Experimental Biology and Ecology* 189: 103-122.

- Cheshire, A.C., Miller, D.J. & Stewart, R., (1999), Effect of dispersed sediment plumes from beach sand replenishment dredging on recruitment of phaeophycean algae to rocky reefs in Gulf St. Vincent, South Australia. The Impact of Sand Dredging on Benthic Community Structure at the Port Stanvac Dredge Site 2: 1-39, University of Adelaide, Adelaide.
- Clarke, K.R. & Warwick, R.M., 2001, Change in Marine Communities: an Approach to Statistical Analysis and Interpretation, PRIMER-E, Plymouth.
- Connell, J.H., Hughes, T.P. & Wallace, C.C., 1997, 'A 30-year study of coral abundance, recruitment, and disturbance at several scales in space and time', *Ecological Monographs* 67: 461-488.
- Denny, M.W., 1983, 'A simple device for recording the maximum force exerted on intertidal organisms', *Limnology and oceanography*: 1269-1274.
- Diaz, M.C. & Rutzler, K., 2001, 'Sponges: an essential component of Caribbean coral reefs', *Bulletin of Marine Science* 69: 535-546.
- Dodge, R.E. & Vaisnys, J.R., 1977, 'Coral populations and growth patterns: responses to sedimentation and turbidity associated with dredging', *Journal of Marine Research* 35: 715-730.
- Done, T.J., 1982, 'Patterns in the distribution of coral communities across the central Great Barrier Reef', *Coral Reefs* 1: 95-107.
- Ebeling, A., Laur, D. & Rowley, R., 1985, 'Severe storm disturbances and reversal of community structure in a southern California kelp forest', *Marine Biology* 84: 287-294.
- Edwards, R.A. & Smith, D.A., 2005, 'Subtidal assemblages associated with a geotextile reef in south-east Queensland, Australia', *Marine and Freshwater Research* 56: 133 142.
- Fellegara, I., 2008, Ecophysiology of the marginal, high-latitude corals (Coelenterata: Scleractinia) of Moreton Bay, QLD, report prepared for Ph.D Thesis, University of Queensland.
- Fisheries Research Consultants, 1991, Survey of the Intertidal and Subtidal Rocky Reefs of the Sunshine Coast, report prepared for Queensland National Parks and Wildlife Service.

- Fisheries Research Consultants, 1995a, Tweed River Entrance Sand Bypassing Project Monitoring Study of the Impacts of Stage 1 (A) Nourishment on Kirra Reef, June 1995, report prepared for Tweed River Entrance Sand Bypassing Project, New South Wales Public Works.
- Fisheries Research Consultants, 1995b, Tweed River Entrance Sand Bypassing Project Monitoring Study of the Impacts of Stage 1 (A) Nourishment on Kirra Reef, May, 1995, report prepared for Tweed River Entrance Sand Bypassing Project, New South Wales Public Works.
- Fisheries Research Consultants, 1996, Tweed River Entrance Sand Bypassing Project, Monitoring Study of the Impacts of Stage 1(A) Nourishment on Kirra Reef, report prepared for Tweed River Entrance Sand Bypassing Project, New South Wales Public Works.
- frc environmental, 2001, Tweed River Entrance Sand Bypassing Project, Monitoring of Kirra Reef, January 2001, report prepared for Brown and Root.
- frc environmental, 2003, *Tweed River Entrance Sand Bypassing Project, Kirra Reef Ecological Monitoring 2003*, report prepared for Department of Infrastructure Planning & Natural Resources (unpublished).
- frc environmental, 2004, *Tweed River Entrance Sand Bypassing Project, Kirra Reef Ecological Monitoring 2004*, report prepared for Department of Infrastructure Planning & Natural Resources (unpublished).
- frc environmental, 2005, Tweed River Entrance Sand Bypassing Project, Kirra Reef Ecological Monitoring Report 2005, report prepared for Department Lands.
- frc environmental, 2010, Tweed River Entrance Sand Bypassing Project, Kirra Reef Ecological Monitoring 2010, report prepared for NSW Land & Property Management Authority and the QLD Department of Environment and Resource Management.
- frc environmental, 2012, *Kirra Reef Marine Biota Monitoring*, report prepared for NSW Department of Primary Industries, Catchment and Land Division.
- Friedlander, A., Brown, E., Jokiel, P., Smith, W. & Rodgers, K., 2003, 'Effects of habitat, wave exposure, and marine protected area status on coral reef fish assemblages in the Hawaiian archipelago', *Coral Reefs* 22: 291-305.

- Harriott, V.J., Banks, S.A., Mau, R.L., Richardson, D. & Roberts, L.G., 1999, 'Ecological and conservation significance of the subtidal rocky reef communities of northern New South Wales, Australia', *Marine and Freshwater Research* 50: 299 306.
- Hatcher, B., Johannes, R. & Robertson, A., 1989, 'Review of research relevant to the conservation of shallow tropical marine ecosystems.', *Oceanography and Marine Biology: an Annual Review.* 27: 337-414.
- Hoey, A. & Bellwood, D., 2008, 'Cross-shelf variation in the role of parrotfishes on the Great Barrier Reef', *Coral Reefs*: 37-47.
- Hollingsworth, P., 1975, 'Environmental Investigation Report: Crown Reserve R68 Currumbin Creek, City of the Gold Coast, Land Administration Commission, unpublished report'.
- Hughes, T.P., 1994, 'Catastrophes, phase shifts, and large scale degradation of a Caribbean coral reef', *Science* 265: 1547-1551.
- Hughes, T.P., Rodrigues, M.J., Bellwood, D.R., Ceccarelli, D., Hoegh-Guldberg, O., McCook, L., Moltschaniwskvi, N., Pratchett, M.S., Steneck, R.S. & Willis, B., 2007, 'Phase shifts, herbivory and the resilience of coral reefs to climate change', *Current Biology* 17: 360-365.
- Hyder Consulting Pty Ltd, Patterson Britton & Partners Pty Ltd & WBM Oceanics Australia Joint Venture, 1997. Tweed River Entrance Sand Bypassing Project, Permanent Bypassing System, Environmental Impact Statement Impact Assessment Study, report no. 9706236-4D June.
- Jompa, J. & McCook, L.J., 2002, 'Effects of competition and herbivory on interactions between a hard coral and a brown alga', *Journal of Experimental Marine Biology and Ecology* 271: 25-39.
- Jones, G.P. & Syms, C., 1998, 'Disturbance, habitat structure and the ecology of fishes on coral reefs', *Australian Journal of Ecology* 23: 287-297.
- Kaufman, L.S., 1983, 'Effects of Hurricane Allen on reef fish assemblages', *Coral Reefs* 2: 43-47.
- Kay, A.M. & Keough, M.J., 1981, 'Occupation of patches in the epifaunal communities on pier pilings and the bivalve Pinna bicolor at Edithburgh, South Australia', *Oecologia* 48: 123-130.

- Kennelly, S., 1987a, 'Inhibition of kelp recruitment by turfing algae and consequences for an Australian kelp community', 112: 49-60.
- Kennelly, S., 1987b, 'Physical disturbances in an Australian kelp community I. temporal effects', *Marine Ecology Progress Series* 40: 145-153.
- Kennelly, S., 1989, 'Effects of kelp canopies on understory species due to shade and scour', *Marine Ecology Progress Series* 50: 215-224.
- Lassig, B.R., 1983, 'The effects of a cyclonic storm on coral reef fish assemblages', Environmental Biology of Fishes 9: 55-63.
- Lesser, M.P., 2006, 'Benthic-pelagic coupling on coral reefs: feeding and growth of Caribbean sponges', *Journal of Experimental Marine Biology & Ecology* 328: 277-288.
- Lopez-Victoria, M., Zea, S. & Wei, E., 2006, 'Competition for space between encrusting excavating Caribbean sponges and other coral reef organisms', *Marine Ecology Progress Series* 312: 113-121.
- Lugo, A.E., Rogers, C. & Nixon, S., 2000, 'Hurricanes, coral reefs and rainforests: resistance, ruin and recovery in the Caribbean', *Ambio* 29: 106-114.
- Massel, S.R. & Done, T.J., 1993, 'Effects of cyclone waves on massive coral assemblages on the Great Barrier Reef: meteorology, hydrodynamics and demography', *Coral Reefs* 12: 153-166.
- McCook, L., Jompa, J. & Diaz-Pulido, G., 2001, 'Competition between corals and algae on coral reefs: a review of evidence and mechanisms', *Coral Reefs* 19: 400-417.
- McCook, L.J., 1997, 'Effects of herbivory on zonation of *Sargassum* spp. within fringing reefs of the central Great Barrier Reef', *Marine Biology* 129: 713-722.
- McGuinness, K.A., 1987, 'Disturbance and organisms on boulders .1. patterns in the environment and the community', *Oecologia* 71: 409-419.
- Miller, R.J. & Etter, R.J., 2008, 'Shading facilitates sessile invertebrate dominance in the rocky subtidal Gulf of Maine', *Ecology* 89: 452-462.
- Moran, D.P. & Reaka Kudla, M.L., 1991, 'Effects of disturbance: disruption and enhancement of coral reef cryptofaunal populations by hurricanes', *Coral Reefs* 9: 215-224.

- Mumby, P., 2009, 'Phase shifts and the stability of macroalgal communities on Caribbean coral reefs', *Coral Reefs* 28: 761-773.
- Murphy, H.M. & Jenkins, G.P., 2010, 'Observational methods used in marine spatial monitoring of fishes and associated habitats: a review', *Marine and Freshwater Research* 61: 236-252.
- Paine, R. & Levin, S., 1981, 'Intertidal Landscapes: Disturbance and the Dynamics of Pattern', *Ecological Monographs* 51: 145-178.
- Pawlik, J.R., Steindler, L., Henkel, T.P., Beer, S. & Ilan, M., 2007, 'Chemical warfare on coral reefs: sponge metabolites differentially affect coral symbiosis in situ', *Limnology and Oceanography* 52: 907-911.
- Reef Check, 2010, http://www.reefcheckaustralia.org/data.html, accessed May 2014.
- Rogers, C.S., 1990, 'Responses of coral reefs and reef organisms to sedimentation', *Marine Ecology Progress Series* 62: 185-202.
- Russ, G.R., 2003, 'Grazer biomass correlates more strongly with production than with biomass of algal turfs on a coral reef', *Coral Reefs* 22: 63-67.
- Rutzler, K., 2002, 'Impact of crustose clionid sponges on Caribbean reef corals', *Acta Geologica Hispanica* 37: 61-72.
- Scheffers, S.R., Nieuwland, G., Bak, R.P.M. & van Duyl, F.C., 2004, 'Removal of bacteria and nutrient dynamics within the coral reef framework of Curacao (Netherlands Antilles)', *Coral Reefs* 23: 413-422.
- Schlacher-Hoenlinger, M., Walker, S., Johnson, J., Schlacher, T. & Hooper, J., 2006, Biological Baseline Survey of the Ex-HMAS Brisbane Artificial Reef, report prepared for Environmental Protection Agency (EPA).
- Schlacher-Hoenlinger, M., Walker, S., Johnson, J., Schlacher, T. & Hooper, J., 2009, Biological Monitoring of the Ex-HMAS Brisbane Artificial Reef: Phase II - Habitat Values, report prepared for Environmental Protection Agency (EPA).
- Shanks, A.L. & Wright, W.G., 1986, 'Adding teeth to wave action: the destructive effects of wave-born rocks on intertidal organisms', *Oecologia* 69: 420-4281.
- Smith, K.A., 2003, 'A simple multivariate technique to improve the design of a sampling strategy for age-based fishery monitoring', *Fisheries Research* 64: 79-85.

- Walker, S., Schlacher, T. & Schlacher-Hoenlinger, M., 2007, 'Spatial heterogeneity of epibenthos on artificial reefs: fouling communities in the early stages of colonization on an East Australian shipwreck', *Marine Ecology* 28: 435-445.
- Walker, S.J., Degnan, B.M., Hooper, J.N.A. & Skilleter, G.A., 2008, 'Will increased storm intensity from climate change affect biodiversity in non-scleractinian sessile fauna on coral reefs?', *Global Change Biology* 14: 2755-2770.
- Warwick, R.M. & Clarke, K.R., 1993, 'Increased variability as a symptom of stress in marine communities', *Journal of Experimental Marine Biology and Ecology* 172: 215-216.
- Woodley, J.D., Chornesky, E.A. & Clifford, P.A., 1981, 'Hurricane Allens impact on Jamaican coral reefs', *Science* 214: 749-755.
- Wulff, J., 2001, 'Assessing and monitoring coral reef sponges: why and how?', *Bulletin of Marine Science* 69: 831-846.

Appendix A Introduction to Data Analysis Used

Multivariate Analyses

Multivariate statistical techniques are widely used in ecology to assess the similarities / relationships between assemblages. Whereas univariate analyses can only compare one variable at a time (e.g. an index of community structure such as a diversity index, or a single indicator species), multivariate analyses can compare samples based on the extent that assemblages share particular taxa and abundances (Clarke & Warwick 2001).

The first step of multivariate analysis usually involves the creation of a matrix of similarity coefficients, computed between every pair of samples. The coefficient is usually a measure of how close the abundance is for each species (defined so that 100% = total similarity and 0% = complete dissimilarity). The Bray Curtis similarity measure is commonly the most appropriate for biological data (Clarke & Warwick 2001).

Multi-dimensional Scaling

Non-metric multi-dimensional scaling ordinations (nMDS) attempt to place samples in two dimensional space, so that the rank order of the distances between samples matches the rank order of the matching similarities from the similarity matrix (Clarke & Warwick 2001). This provides a visual representation of the similarities between assemblages within each sample. Each of the axes is not related to any particular value; in fact axes can be rotated to provide the best visual representation of the data. Ordinations are particularly useful tools for analysing, and visually presenting, differences between assemblages. Ordinations are essentially maps of samples, in which the placement of samples on the map reflects the similarly of the community to the communities in other samples (Clarke & Warwick 2001). Distances between samples on an ordination attempt to match the similarities in assemblage structure: nearby points represent assemblages with very few attributes (species or abundance of species); points far apart have very few attributes in common (Clarke & Warwick 2001).

A stress coefficient is calculated to reflect the extent to which the multi-dimensional scaling ordination and the similarity matrix agree (Clarke & Warwick 2001) (i.e. how well the multi-dimensional scaling ordination accurately reflects the relationship between samples). Stress values of <0.15 are generally acceptable.

In Figure A2, each freshwater macroinvertebrate sample is represented on the multi-dimensional scaling ordination. By looking at the distances between each sample,

we can infer that samples (assemblages) from the same stream reach (e.g. sites DS, M, STC and US) group together. That is, they are more similar to each other than they are to samples taken from other stream reaches.




Figure A2 Example of a multi-dimensional scaling ordination for macroinvertebrate communities sampled in riffle habitats of different stream reaches.

Analysis of Similarity

ANOSIM is analogous to ANOVA in univariate statistics (Smith 2003). A global R statistic is calculated to determine whether there is a significant difference between all samples. If there are differences, then pairwise comparisons are conducted to test for differences between pairs of samples (analogous to post-hoc tests in ANOVA).

The R value lies between -1 and +1 (all similarities within groups are less than any similarity between groups), with a value of zero representing the null hypothesis (no difference among a set of samples) (Clarke & Warwick 2001). Comparison of pairwise R values can give an indication of how different assemblages are: R values close to 0 indicate little difference, values around 0.5 indicate some overlap and values close to 1 to indicate many or substantial differences. In many instances however, researches are primarily interested in whether the R value is statistically different from zero (usually at a confidence level of 0.05) (Clarke & Warwick 2001) (i.e. whether they can reject the null hypothesis).

ANOSIM can provide information on whether the (visual) differences between assemblages in the multi-dimensional scaling ordination are significant; it is an independent test from the multi-dimensional scaling ordination. It is based on testing the

differences between the rank similarities in the similarity matrix, not on the distances between samples in the multi-dimensional scaling ordination (Clarke & Warwick 2001).

Permutational Multivariate Analysis of Variance

PERMANOVA is used to test simultaneous responses of one or more variables to one or more factors in an *a priori* structured design, using random permutation of the data to assess significance (Anderson 2004). PERMANOVA generates a pseudo F-statistic similar to traditional ANOVA, but p-values are calculated with permutations, which does not assume normal data distribution. PERMANOVA can provide information on whether the (visual) differences between assemblages in a multi-dimensional scaling ordination are significant; however, it is an independent test from the multi-dimensional scaling ordination.

Were significant differences among factors are found, post-hoc pairwise comparison can then be used to test for differences between pairs of samples (analogous to post-hoc tests in ANOVA).

The level of multivariate dispersion among samples within each of the test groups can be examined using the permutational analysis of multivariate dispersions (PERMDISP) routine (Anderson 2004). In traditional impact assessment, a change in the dispersion of data can also indicate an impact.

Similarity Percentage – Species Contributions

SIMPER analysis provides information on how dissimilar assemblages from various groups are (e.g. how similar all of the macroinvertebrate samples taken for a particular habitat within a stream reach are), and how similar each group (e.g. reach) is to any other group. SIMPER analysis also identifies the species / taxa that are contributing to the dissimilarity between two assemblages, in rank order (i.e. it identifies which species is contributing the most to the differences). SIMPER analysis may help to identify potential 'indicator' species. For example, if a particular species consistently contributes greatly to the differences between impacted and unimpacted assemblages, it may be a useful indicator. The abundance of this indicator species can then be compared between sites using univariate techniques such as ANOVA.

Appendix B Cover and Abundance of Benthic Fauna and Flora on Kirra and Palm Beach Reefs in April 2014

Table B. 1 R statistic of pairwise ANOSIM results for differences in the composition of benthic assemblages between surveys for each reef.

Reef	1995-1996	1996-2001	2001-2003	2003-2004	2004-2005	2005-2010	2010-2012	2012-2014
Kirra Reef	0.412	0.101	0.280	0.226	0.468	0.130	0.665	0.598
Palm Beach Reef	0.080	0.516	0.413	0.296	0.182	0.832	0.507	0.114

Table B.2 Benthic assemblage data from Kirra and Palm Beach reefs in April 2014.

Location	Site	Replicate	% Macroalgae	% Turf algae	% Soft coral	% Hard coral	% Sponge	% Ascidians	% Bare	% Anemone	% Barnacle	% Crustose coralline algae	# Crinoids	# <i>Pyura</i> sp.	# Echinoidea	# Tube worms	# Polychaetes	# Cnemidocarpa sp.
Palm Beach	1	1	0	20	0	80	0	0	0	0	0	0	0	0	0	0	0	0
Palm Beach	1	2	0	30	20	0	40	5	5	0	0	0	0	1	0	0	0	0
Palm Beach	1	3	0	0	40	0	0	0	50	0	10	0	0	0	0	0	0	0
Palm Beach	1	4	0	20	0	70	0	5	5	0	0	0	0	0	0	0	0	0
Palm Beach	1	5	0	30	10	50	0	0	10	0	0	0	0	0	0	0	0	0
Palm Beach	1	6	0	25	30	5	0	0	0	0	40	0	0	0	0	0	0	0

Location	Site	Replicate	% Macroalgae	% Turf algae	% Soft coral	% Hard coral	% Sponge	% Ascidians	% Bare	% Anemone	% Barnacle	% Crustose coralline algae	# Crinoids	# <i>Pyura</i> sp.	# Echinoidea	# Tube worms	# Polychaetes	# Cnemidocarpa sp.
Palm Beach	1	7	0	30	20	30	0	0	0	0	20	0	0	0	0	0	0	0
Palm Beach	1	8	0	30	55	0	0	0	0	5	10	0	0	0	0	0	0	0
Palm Beach	1	9	0	50	20	15	0	5	0	0	10	0	0	0	0	0	0	1
Palm Beach	1	10	0	45	40	0	0	5	0	10	0	0	0	0	0	0	0	1
Palm Beach	1	11	0	40	5	40	15	0	0	0	0	0	0	0	0	0	0	0
Palm Beach	1	12	5	40	20	10	0	5	0	20	0	0	0	2	0	0	0	1
Palm Beach	1	13	0	50	40	0	0	10	0	0	0	0	0	1	0	0	0	1
Palm Beach	1	14	0	60	0	0	30	0	10	0	0	0	0	0	0	0	0	0
Palm Beach	1	15	0	30	30	0	40	0	0	0	0	0	0	0	0	0	0	0
Palm Beach	2	1	0	0	0	0	0	80	20	0	0	0	0	9	0	0	0	0

Location	Site	Replicate	% Macroalgae	% Turf algae	% Soft coral	% Hard coral	% Sponge	% Ascidians	% Bare	% Anemone	% Barnacle	% Crustose coralline algae	# Crinoids	# Pyura sp.	# Echinoidea	# Tube worms	# Polychaetes	# Cnemidocarpa sp.
Palm Beach	2	2	0	0	0	0	5	70	15	0	10	0	0	12	0	0	0	0
Palm Beach	2	3	0	0	20	20	0	20	30	0	10	0	0	7	0	0	0	0
Palm Beach	2	4	0	0	90	0	0	0	0	0	10	0	0	0	0	0	0	0
Palm Beach	2	5	0	0	0	40	0	5	5	0	50	0	0	0	0	0	0	0
Palm Beach	2	6	0	0	0	0	25	5	10	0	60	0	0	2	0	0	0	0
Palm Beach	2	7	0	0	5	20	10	10	10	0	45	0	0	0	0	0	0	0
Palm Beach	2	8	0	0	0	30	0	5	45	0	20	0	0	0	0	5	0	0
Palm Beach	2	9	0	20	0	80	0	0	0	0	0	0	0	0	0	0	0	0
Palm Beach	2	10	5	15	40	40	0	0	0	0	0	0	0	0	0	0	0	0
Palm Beach	2	11	0	30	50	0	10	0	10	0	0	0	0	0	0	0	0	0

Location	Site	Replicate	% Macroalgae	% Turf algae	% Soft coral	% Hard coral	% Sponge	% Ascidians	% Bare	% Anemone	% Barnacle	% Crustose coralline algae	# Crinoids	# <i>Pyura</i> sp.	# Echinoidea	# Tube worms	# Polychaetes	# Cnemidocarpa sp.
Palm Beach	2	12	0	70	0	20	10	0	0	0	0	0	0	0	0	0	0	0
Palm Beach	2	13	0	25	40	25	0	0	0	0	10	0	0	0	0	0	0	0
Palm Beach	2	14	20	25	5	20	0	10	0	0	20	0	0	0	0	0	0	1
Palm Beach	2	15	0	30	0	25	0	5	0	0	40	0	0	0	0	0	0	0
Palm Beach	3	1	0	60	0	5	0	1	19	0	15	0	0	0	0	0	0	0
Palm Beach	3	2	0	70	0	10	0	5	5	0	10	0	0	0	0	0	0	1
Palm Beach	3	3	0	61	16	1	1	5	0	0	16	0	0	0	0	0	0	0
Palm Beach	3	4	0	79	0	1	10	5	0	0	5	0	0	0	0	0	0	0
Palm Beach	3	5	0	35	20	10	5	0	25	0	5	0	0	0	0	0	0	0
Palm Beach	3	6	0	55	15	5	5	0	10	0	10	0	0	0	0	0	0	0

Location	Site	Replicate	% Macroalgae	% Turf algae	% Soft coral	% Hard coral	% Sponge	% Ascidians	% Bare	% Anemone	% Barnacle	% Crustose coralline algae	# Crinoids	# <i>Pyura</i> sp.	# Echinoidea	# Tube worms	# Polychaetes	# Cnemidocarpa sp.
Palm Beach	3	7	0	20	0	0	20	60	0	0	0	0	0	0	0	0	3	0
Palm Beach	3	8	0	40	0	10	20	5	0	20	5	0	0	0	0	1	0	0
Palm Beach	3	9	0	65	0	0	0	30	0	0	5	0	0	0	0	0	1	0
Palm Beach	3	10	20	40	0	0	40	0	0	0	0	0	0	0	0	0	0	0
Palm Beach	3	11	0	50	0	30	10	10	0	0	0	0	0	1	0	0	0	1
Palm Beach	3	12	0	30	30	40	0	0	0	0	0	0	0	0	0	0	4	0
Palm Beach	3	13	0	40	20	40	0	0	0	0	0	0	0	0	1	0	0	0
Palm Beach	3	14	0	60	0	10	0	30	0	0	0	0	0	0	0	0	0	0
Palm Beach	3	15	0	50	0	0	20	0	0	30	0	0	0	0	1	0	0	0
Kirra Reef	KRO 3	1	0	20	0	0	0	60	20	0	0	0	0	0	0	0	0	0

Location	Site	Replicate	% Macroalgae	% Turf algae	% Soft coral	% Hard coral	% Sponge	% Ascidians	% Bare	% Anemone	% Barnacle	% Crustose coralline algae	# Crinoids	# <i>Pyura</i> sp.	# Echinoidea	# Tube worms	# Polychaetes	# Cnemidocarpa sp.
Kirra Reef	KRO 3	2	19	38	0	0	5	38	0	0	0	0	0	3	0	0	0	0
Kirra Reef	KRO 3	3	40	0	0	0	5	20	35	0	0	0	0	2	0	0	0	0
Kirra Reef	KRO 3	4	5	20	0	0	10	60	5	0	0	0	0	2	0	0	0	8
Kirra Reef	KRO 3	5	30	40	0	0	0	10	20	0	0	0	0	2	0	0	0	0
Kirra Reef	KRO 3	6	40	20	0	0	0	40	0	0	0	0	0	1	0	0	0	6
Kirra Reef	KRO 3	7	10	30	0	0	20	40	0	0	0	0	0	0	0	0	0	5
Kirra Reef	KRO 3	8	30	30	0	0	0	30	10	0	0	0	0	3	0	0	0	1
Kirra Reef	KRO 3	9	10	50	0	0	15	20	0	0	0	5	1	0	0	0	0	2
Kirra Reef	KRO 3	10	30	20	0	0	20	20	0	0	0	10	0	2	0	0	0	0
Kirra Reef	KRO 3	11	20	60	0	0	0	10	10	0	0	0	0	1	0	0	0	0

Location	Site	Replicate	% Macroalgae	% Turf algae	% Soft coral	% Hard coral	% Sponge	% Ascidians	% Bare	% Anemone	% Barnacle	% Crustose coralline algae	# Crinoids	# <i>Pyura</i> sp.	# Echinoidea	# Tube worms	# Polychaetes	# Cnemidocarpa sp.
Kirra Reef	KRO 3	12	30	20	0	0	10	30	10	0	0	0	0	2	0	1	0	3
Kirra Reef	KRO 3	13	30	30	0	0	10	30	0	0	0	0	0	2	0	0	0	2
Kirra Reef	KRO 3	14	20	30	0	0	0	50	0	0	0	0	3	1	0	0	0	2
Kirra Reef	KRO 3	15	20	30	0	0	10	20	20	0	0	0	0	2	0	0	0	4
Kirra Reef	KRN1	1	30	47	0	0	0	1	0	0	2	20	0	1	0	0	0	0
Kirra Reef	KRN1	2	40	33	0	0	0	5	10	0	2	10	0	2	0	0	0	0
Kirra Reef	KRN1	3	30	35	0	0	0	5	0	0	0	30	0	3	0	0	0	0
Kirra Reef	KRN1	4	20	70	0	0	0	0	0	0	0	10	0	0	0	0	0	0
Kirra Reef	KRN1	5	40	39	0	0	5	5	0	0	1	10	0	0	0	0	0	0
Kirra Reef	KRN1	6	40	20	0	0	0	0	0	0	30	10	0	0	0	0	0	3

Location	Site	Replicate	% Macroalgae	% Turf algae	% Soft coral	% Hard coral	% Sponge	% Ascidians	% Bare	% Anemone	% Barnacle	% Crustose coralline algae	# Crinoids	# Pyura sp.	# Echinoidea	# Tube worms	# Polychaetes	# Cnemidocarpa sp.
Kirra Reef	KRN1	7	20	45	0	0	0	0	0	0	15	20	0	0	0	0	0	0
Kirra Reef	KRN1	8	35	60	0	0	0	5	0	0	0	0	0	2	0	0	0	0
Kirra Reef	KRN1	9	20	64	0	0	1	10	0	0	0	5	2	0	0	0	0	0
Kirra Reef	KRN1	10	20	70	0	0	0	5	0	0	0	5	0	3	0	0	0	0
Kirra Reef	KRN1	11	30	54	0	0	0	5	0	0	1	10	1	3	0	0	0	0
Kirra Reef	KRN1	12	50	29	0	0	5	15	0	0	1	0	0	3	0	1	0	0
Kirra Reef	KRN1	13	30	20	0	0	10	20	0	0	0	20	0	4	0	1	0	0
Kirra Reef	KRN1	14	10	64	0	0	0	15	1	0	0	10	0	5	0	0	0	0
Kirra Reef	KRN1	15	10	50	0	0	0	40	0	0	0	0	0	0	0	0	0	0
Kirra Reef	KRN2	1	30	55	0	0	5	5	0	0	0	5	0	1	0	0	0	0

Location	Site	Replicate	% Macroalgae	% Turf algae	% Soft coral	% Hard coral	% Sponge	% Ascidians	% Bare	% Anemone	% Barnacle	% Crustose coralline algae	# Crinoids	# <i>Pyura</i> sp.	# Echinoidea	# Tube worms	# Polychaetes	# Cnemidocarpa sp.
Kirra Reef	KRN2	2	20	60	0	0	10	10	0	0	0	0	1	2	0	0	0	0
Kirra Reef	KRN2	3	15	50	0	0	0	20	0	5	0	10	0	3	0	0	0	0
Kirra Reef	KRN2	4	20	10	0	0	0	30	10	10	0	20	0	1	0	0	0	0
Kirra Reef	KRN2	5	20	40	0	0	10	0	0	0	0	30	2	0	0	0	0	1
Kirra Reef	KRN2	6	10	50	0	0	10	20	0	0	0	10	1	0	0	0	0	3
Kirra Reef	KRN2	7	10	50	0	0	10	20	10	0	0	0	0	0	0	0	0	1
Kirra Reef	KRN2	8	5	0	5	0	0	40	15	30	0	5	1	2	0	0	0	0
Kirra Reef	KRN2	9	20	10	0	0	0	40	0	0	0	30	0	0	0	0	0	0
Kirra Reef	KRN2	10	20	10	0	0	10	40	0	20	0	0	3	1	0	0	0	3
Kirra Reef	KRN2	11	60	20	0	0	0	20	0	0	0	0	0	0	0	0	0	1

Location	Site	Replicate	% Macroalgae	% Turf algae	% Soft coral	% Hard coral	% Sponge	% Ascidians	% Bare	% Anemone	% Barnacle	% Crustose coralline algae	# Crinoids	# Pyura sp.	# Echinoidea	# Tube worms	# Polychaetes	# Cnemidocarpa sp.
Kirra Reef	KRN2	12	50	20	0	0	0	10	0	0	0	20	0	2	0	0	0	0
Kirra Reef	KRN2	13	30	20	0	0	0	0	20	15	0	15	1	0	0	0	0	0
Kirra Reef	KRN2	14	30	35	0	0	15	15	0	5	0	0	0	1	0	0	0	2
Kirra Reef	KRN2	15	20	50	0	0	0	30	0	0	0	0	2	4	0	0	0	1

Appendix C Relative Abundance of Fish found at Kirra and Palm Beach Reefs in Each Survey

	Table C1	Fish species and their relative abundance in April 2014 and in previous surv	eys.
--	----------	--	------

	0					Kirra	Reef									Palm Be	ach Reef				
Scientific Name	Common Name	Apr '95	Jun '95	Feb '96	Jan '01	May '03	Mar '04	Feb '05	Feb '10	Jul '12	Apr '14	Apr '95	Jun '95	Feb '96	Jan '01	Мау '03	Mar '04	Feb '05	Feb '10	Jul '12	Apr '14
Acanthuridae																					
Acanthurus grammoptilus	ring-tailed surgeon	**	**	**	**	**		**	**		***	**	**	***	**	**		**	**	**	**
Acanthurus xanthopterus	yellowfin surgeonfish									**	**									**	
Prionurus microlepidotus	sawtail surgeon								***									**	**	**	
Apogonidae																					
Apogon cookii	Cook's cardinal fish															_	**	**	*		
Apogon doederleini	four lined cardinal fish							**	*		*				**		***	**	**		
Aracanidae																					
Strophiurichthys robustus	freckled boxfish					**					*										
Balastidae																					
Sufflamen chrysopterus	half-moon triggerfish							**			**				_			**	**	*	**
Sufflamen fraenatus	bridled triggerfish													*							
Blenniidae																					
Plagiotremus tapeinosoma	hit and run blenny						**	*	*	*	*							*	**		**
Brachaeluridae																					
Brachaelurus waddi	blindshark							*		*							*			*	*
Carangidae																					
Caranx sp.	trevally										*		**	***	***						
Gnathanodon speciosus	golden trevally																**				

	0					Kirra	Reef									Palm Be	ach Reef				
Scientific Name	Common Name	Apr '95	Jun '95	Feb '96	Jan '01	May '03	Mar '04	Feb '05	Feb '10	Jul '12	Apr '14	Apr '95	Jun '95	Feb '96	Jan '01	May '03	Mar '04	Feb '05	Feb '10	Jul '12	Apr '14
Pseudocaranx dentex	silver trevally								***							***					
Trachinotus blochii	dart	**									*										
Trachurus novaezelandiae	yellowtail	****	****	***		****	***	****	****	***	***	****	****	**	****	****			***	***	
Chaetodontidae																					
Chaetodon auriga	threadfin butterfly fish	**	**			**			**		*	*	*	**			**				**
Chaetodon citrinellus	citron butterfly fish					**												**	**	**	
Chaetodon flavirostris	dusky butterfly fish				*						*							**	**		**
Chaetodon lineolatus	lined butterfly fish				*	*					*			**				*			**
Henochus sp.	banner fish	**				*			**												
Cheilodactylidae	,																				
Cheilodactylus fuscus	red morwong			**	**	**	**	*	**	*	**		**		**	**		**	**		
Cheilodactylus vestitus	crested morwong			*				*	*	*	**				*			**	**		*
Chironemidae																					
Chironemus marmoratus	kelp fish				**	**		**			**										
Cirrhitidae																					
Cirrhitichthys sp.	hawkfish					**								*			**	**			*
Dasyatidae																					
Dasyatis kuhlii	blue-spotted maskray								**												
Pastinachus atrus	cowtail stingray										*										
Diodontidae																					
Dicotylichthys punctulatus	three-bar porcupine fish					**	*	***	**	**	**				*						**

	Commen					Kirra	Reef							·		Palm Be	ach Reef				
Scientific Name	Common Name	Apr '95	Jun '95	Feb '96	Jan '01	May '03	Mar '04	Feb '05	Feb '10	Jul '12	Apr '14	Apr '95	Jun '95	Feb '96	Jan '01	Мау '03	Mar '04	Feb '05	Feb '10	Jul '12	Apr '14
Diodon holocanthus	freckled porcupine fish	*		**																**	
Diodon hystrix	black- spotted porcupine fish					*															
Ephippidae																					
Platax orbicularis	round batfish								*												
Enoplosidae																					
Enoplosus armatus	old wife	***		**	***			**			*				**						
Fistularidae																					
Fistularia commersonii	smooth flutemouth	*			**				**										**		
Fistularia petimba	rough flutemouth																		**		
Gerreidae																					
Gerres subfasciatus	silver biddy	***	**	**	**	***	*	**	*		**										
Haemulidae																					
Plectorhinchus flavomaculatus	gold-spotted sweetlip				*	**		*			*							*	*		*
Labridae																					
Achoerodus viridis	blue groper															*					
Anampses meleagrides	spotted wrasse																			*	
Diproctacanthus xanthurus	yellowtail tubelip									*											
Halichoeres sp.	striped wrasse			**	***	*			***	***	***				**		*	**	**		***
Labroides dimidiatus	cleaner wrasse			**	**	***	**	**	**	**	**		*	*	*	**	**	*	**	**	***

	0					Kirra	Reef									Palm Be	ach Reef				
Scientific Name	Common Name	Apr '95	Jun '95	Feb '96	Jan '01	May '03	Mar '04	Feb '05	Feb '10	Jul '12	Apr '14	Apr '95	Jun '95	Feb '96	Jan '01	May '03	Mar '04	Feb '05	Feb '10	Jul '12	Apr '14
Notolabrus gymnogenis	crimson- banded wrasse					***	**													*	
Notolabrus sp.	wrasse	**										**	**	**							
Pseudolabrus guentheri	Gunther's wrasse									**	***									**	**
Thalassoma jansenii	Jansen's wrasse									**										*	*
Thalassoma lunare	moon wrasse					**	**	**		**	**			***	***	***	**	***	***	**	***
Thalassoma lutescens	yellow moon wrasse	**		**	**	***	*	**		**	*		**	***	***	***	**	**	***		***
Lutjanidae																					
Lutjanus fulviflamma	black-spot snapper									**	*										
Microcanthidae																					
Atypichthys strigatus	Australian mado			**	***	***	*	***	***		***						***	***	***	**	
Microcanthus strigatus	stripey	***	***	**	***	***	**	***	***	**	*	***	**	**	***		*		***	**	***
Monacanthidae																					
Meuschenia trachylepis	yellow-tailed leatherjacket					*												**			
Monacanthus chinensis	fan-bellied leatherjacket	*		*	*				*				*	*				**			
Oxymonacanthus Iongirostris	orange spotted filefish					-				_										*	*
Monodactylidae																					
Monodactylus argenteus	silver batfish					***							***		**	***					
Schuettea scalaripinnis	eastern pomfred								***		*	***	***	***			_		**		

	Common					Kirra	Reef									Palm Be	ach Reef				
Scientific Name	Common Name	Apr '95	Jun '95	Feb '96	Jan '01	May '03	Mar '04	Feb '05	Feb '10	Jul '12	Apr '14	Apr '95	Jun '95	Feb '96	Jan '01	May '03	Mar '04	Feb '05	Feb '10	Jul '12	Ap '14
Mullidae																					
Parupeneus barberinoides	half-and-half goatfish				**			*										**			
Parupeneus ciliatus	diamond- scaled goat fish										*			*		_					
Parupeneus signatus	black spot goat fish	***	**	***	***										***					**	*
Muraenidae																					
Gymnothorax favagineus	tessellate moray					*	*				*			*						*	*
Gymnothorax prasinus	green moray					**	**	***											*		*
Siderea thyrsoidea	white-eyed moray						**	**													
Myliobatididae																					
Aetobatus ocellatus	white- spotted eagle ray									**	*										
Orectolobidae																					
Orectolobus ornatus	ornate wobbegong					**	**	**	**	**	*		**	**		**		**		**	
Ostraciidae					'																
Ostracion cubicus																		**			
Pempheridae																					
Pempheris multiradiata	bullseye									**			***	***	**				**		*
Pempheris oualensis	black-finned bullseye					*												,			
Pinguipedidae																					
Parapercis queenslandiae	Queensland grubfish																				*

	Common	Kirra Reef												Palm Beach Reef										
Scientific Name	Common Name	Apr '95	Jun '95	Feb '96	Jan '01	May '03	Mar '04	Feb '05	Feb '10	Jul '12	Apr '14	Apr '95	Jun '95	Feb '96	Jan '01	Мау '03	Mar '04	Feb '05	Feb '10	Jul '12	Apr '14			
Platycephalidae																								
Platycephalus fuscus	dusky flathead				*			*																
Plotosidae																								
Cnidoglanis macrocephala	estuary catfish	*																						
Polynemidae																								
Polydactylus nigripinnis	black-finned threadfin															*								
Pomacanthidae																								
Centropyge tibicen	keyhole angelfish					*		*				'					*	**	**					
Pomacanthus semicirculatus	blue angelfish																		*					
Pomacentridae																								
Abudefduf bengalensis	Bengal sergeant major					**		*								**	**	**	*	**	*			
Abudefduf vaigiensis	sergeant major				,				*		*					***	*	**	***		**			
Abudefduf saxatilis	five-banded sergeant major								**							***			*					
Amphiprion sp.	clown fish	**	**	**		**					*			**		**			***	**	*			
Chromis chrysura	robust puller			**	'						**						*				***			
Chromis nitida	barrier reef chromis			**							**			**							***			
Chrysiptera sp.	demoiselle					**										**	*		**					
Dascyllus trimaculatus	domino puller							*							*						**			
Parma microlepis	white ear puller		**	**	**							**	**	**	**		*	*	*					

	Common					Kirra	Reef			Palm Beach Reef											
Scientific Name	Common Name	Apr '95	Jun '95	Feb '96	Jan '01	May '03	Mar '04	Feb '05	Feb '10	Jul '12	Apr '14	Apr '95	Jun '95	Feb '96	Jan ʻ01	May '03	Mar '04	Feb '05	Feb '10	Jul '12	Ap '14
Parma oligolepis	large-scaled parma					**	**	**		**	***							*	***		*
Parma polylepis	banded parma							**										**	**		
Plectroglyphidodon leucozonus	whiteband damsel				ı															*	
Pomacentrus australis	Australian damsel	**	**	**								**	**			**	**				
Pomacentrus coelestis	neon damsel					***		**			***					***		***	****	**	*
Stegastes gascoynei	coral sea gregory					**	*										**		**		
Stegastes sp.	damsel									*										***	
Pomatomidae																					
Pomatomus saltatrix	tailor	****										'									
Rhinobatidae																					
A <i>ptychotrema</i> sp.	shovelnose ray	*									*										
Glaucostegus typus	giant shovelnose ray																		*		
Scorpaenidiae																					
Centropogon australis	fortescue									**					*						
Pterois volitans	red firefish					*										*					
Scorpaena cardinalis	red scorpionfish					*										*			**		
Synanceia horrida	estuarine stonefish							**													
Scombridae																					
Cybiosarda elegans	leaping bonito								****												

	Kirra Reef												Palm Beach Reef										
Scientific Name	Common Name	Apr '95	Jun '95	Feb '96	Jan '01	May '03	Mar '04	Feb '05	Feb '10	Jul '12	Apr '14	Apr '95	Jun '95	Feb '96	Jan '01	Мау '03	Mar '04	Feb '05	Feb '10	Jul '12	Apr '14		
Scomberomorus commerson	spanish mackerel																		**				
Scorpididae																							
Scorpis lineolatus	sweep	***	***	**	**	**	*	**	**	***	*	***	***	**	****	**	**	**	***	****	*		
Serranidae																							
Anthias sp.	anthias										*										*		
Diploprion bifasciatum	barred soapfish									,											*		
Epinephelus fasciatus	black-tipped cod												**				*	***	**		*		
Plectropomus maculatus	coral trout												*										
Siganidae																_							
Siganus fuscescens	rabbitfish				***					*					***		***	***	***		*		
Sillaginidae																							
Sillago analis	gold-lined whiting							**			*												
Sparidae																							
Acanthopagrus australis	yellow fin	***	***	**	***	***				**	***	***	**	**	**				***	**	**		
Rhabdosargus sarba	tarwhine	***			**						*								*				
Sphyraenidae																							
Sphyraena argentea	barracuda										**												
Sphyraena obtusata	striped sea pike	****	**	**	**	*		****		**	**					*			**	**			
Syngnathidae																							
Signathid sp. 1	pipefish	*																					
Stegostomatidae																							
Stegostoma fasciatum	leopard shark														*								

	0	Kirra Reef														Palm Be	ach Reef				
Scientific Name	Common Name	Apr '95	Jun '95	Feb '96	Jan '01	May '03	Mar '04	Feb '05	Feb '10	Jul '12	Apr '14	Apr '95	Jun '95	Feb '96	Jan '01	May '03	Mar '04	Feb '05	Feb '10	Jul '12	Apr '14
Tetraodontidae																					
Arothron hispidus	stars and stripes pufferfish				*		**	**	*						*	*					*
Arothron immaculatus	immaculate pufferfish			*																	
Arothron manilensis	narrow lined toadfish				*																
Arothron stellatus	starry toadfish							*	*								**	**			
Canthigaster valentini	black- saddled toby							*								**		**			
Lagocephalus sp.	toadfish				***																
Torquigener pleurogramma	toadfish	*	*		***									*							
Urolophidae																					
Urolophus sp.	stingaree	*	***	**	**	*		**			*										
Zanclidae																					
Zanclus cornutus	Moorish idol										*										

shading indicates surveys where each species was observed

^{* &}lt; 5 individuals

^{** 6-20} individuals

^{*** 21-100} individuals

^{**** &}gt;100 individuals